

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau


## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>A61K</b>	<b>A2</b>	(11) International Publication Number: <b>WO 97/11669</b> (43) International Publication Date: 3 April 1997 (03.04.97)
(21) International Application Number: PCT/US96/15346 (22) International Filing Date: 25 September 1996 (25.09.96) (30) Priority Data: 08/533,895 26 September 1995 (26.09.95) US (71) Applicant: THE GOVERNMENT OF THE UNITED STATES OF AMERICA, represented by THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES, Office of Technology Transfer National Institutes of Health [US/US]; Suite 325, 6011 Executive Boulevard, Rockville, MD 20852 (US). (72) Inventors: TOPALIAN, Suzanne, L.; 19400 James Creek Court, Brookeville, MD 20833 (US). ROSENBERG, Steven, A.; 10104 Iron Gate Road, Potomac, MD 20854 (US). ROBBINS, Paul, F.; 13009 Bluhill Road, Silver Spring, MD 20906 (US). (74) Agent: FEILER, William, S.; Morgan & Finnegan, L.L.P., 345 Park Avenue, New York, NY 10154 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>

(54) Title: MHC-CLASS II RESTRICTED MELANOMA ANTIGENS AND THEIR USE IN THERAPEUTIC METHODS

## (57) Abstract

The present invention provides MHC Class II restricted melanoma antigens recognized by CD4<sup>+</sup> T cells. This invention further provides prophylactic and therapeutic applications for the Class II restricted melanoma antigens. In particular, this invention provides tyrosinase Class II restricted melanoma antigens, as well as tyrosinase immunogenic peptides which have been modified to enhance their immunogenicity. These antigens can serve as immunogens or vaccines to prevent or treat melanoma. In addition a method for isolating Class II restricted melanoma antigens or identifying new Class II restricted melanoma antigens is provided.



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

- 1 -

TITLE OF THE INVENTION

MHC-CLASS II RESTRICTED MELANOMA ANTIGENS AND THEIR USE IN  
THERAPEUTIC METHODS

FIELD OF THE INVENTION

5           This invention is in the field of prevention and  
treatment of human cancers. More specifically, this  
invention relates to MHC Class II restricted melanoma  
antigens recognized by helper T Cells and to the  
preventative, and therapeutic applications which employ  
10 these antigens. This invention also relates to methods  
for determining Class II restricted melanoma antigens.

BACKGROUND OF THE INVENTION

          Melanomas are aggressive, frequently metastatic  
tumors derived from either melanocytes or melanocyte  
15 related nevus cells ("Cellular and Molecular Immunology"  
(1991) (eds) Abbas A.K., Lechtman, A.H., Pober, J.S.; W.B.  
Saunders Company, Philadelphia: pages 340-341). Melanomas  
make up approximately three percent of all skin cancers  
and the worldwide increase in melanoma is unsurpassed by  
20 any other neoplasm with the exception of lung cancer in  
women ("Cellular and Molecular Immunology" (1991) (eds)  
Abbas, A.K., Lechtman, A.H., Pober, J.S.; W.B. Saunders  
Company Philadelphia pages: 340-342; Kirkwood and Agarwala  
(1993) Principles and Practice of Oncology 7:1-16). Even  
25 when melanoma is apparently localized to the skin, up to  
30% of the patients will develop systemic metastasis and  
the majority of these will die (Kirkwood and Agarwala  
(1993) Principles and Practice of Oncology 7:1-16).  
Classic modalities of treating melanoma include surgery,  
30 radiation and chemotherapy. In the past decade  
immunotherapy and gene therapy have emerged as new and  
promising methods for treating melanoma (Biologic Therapy  
of Cancer, 2nd ed. (1995) Devita, V.T., Hellman, S, and  
Rosenberg, S.A., eds; J.B. Lippincott Co., Philadelphia).

35

- 2 -

Shared melanoma-associated antigens (Ag) expressed among a variety of melanoma patients can be recognized by cytotoxic CD8<sup>+</sup> T lymphocytes derived from melanoma patients. In short-term lysis assays, cytotoxic T lymphocytes (CTL) grown from in vitro sensitized peripheral blood lymphocytes (PBL) or lymph node lymphocytes, or from lymphocytes infiltrating metastatic melanoma lesions, have been shown to recognize autologous and MHC class I compatible allogeneic melanomas but not HLA-matched nonmelanoma tumors, lymphoblasts, or cultured fibroblasts (Darrow, D.L., Slingluff, C.L., & Siegler, H.F. (1989.) *J. Immunol.* **142**, 3329-3335; Hom, S.S., et al. (1991) *J. Immunother.* **10**, 153-164). Similar recognition patterns have been observed by measuring cytokine secretion from tumor infiltrating lymphocytes (TIL) cocultivated with autologous or HLA-matched allogeneic tumor stimulators (Hom, S.S., et al. (1993) *J. Immunother.* **13**, 18-30). Recently, melanoma-specific HLA-A2 restricted CTL clones have been shown to recognize cultured normal melanocytes as well as their malignant counterparts, suggesting that shared melanoma antigens can be lineage specific (Anichini, A., et al. (1993) *J. Exp. Med.* **177**, 989-998). To date, several class I-restricted melanoma-associated antigens have been molecularly defined (Van Der Bruggen, P., et al. (1991) *Science* **254**, 1643-1647; Brichard, V., et al. (1993) *J. Exp. Med.* **178**, 489-495; Kawakami, Y., et al. (1994) *Proc. Natl. Acad. Sci. USA.* **91**, 3515-3519; Bakker, A.B.H., et al. (1994) *J. Exp. Med.* **179**, 1005-1009; Kawakami, Y., et al. (1994) *Proc. Natl. Acad. Sci. USA.* **91**, 6458-6462; Gaugler B., et al. (1994) *J. Exp. Med.* **179**, 921-930). These antigens and derivative class I-restricted peptides 8 to 10 amino acids in length are currently being developed as clinical vaccines to stimulate CD8<sup>+</sup> T cell responses against melanoma.

While animal models of malignant and viral diseases have shown the importance of CD8<sup>+</sup> T cells in the

- 3 -

effector phase of the immune response, the CD4<sup>+</sup> helper arm has been shown to mediate critical priming and effector functions as well. T cell receptors on CD4<sup>+</sup> T cells recognize a complex consisting of an antigenic peptide in conjunction with an MHC Class II molecule. Unlike peptides binding to MHC class I molecules, which are restricted in length from 8-10 amino acids, the antigenic peptides that bind Class II range from about 10 to about 34 amino acids in length and even entire proteins. (Chicz, R.M. et al (1993) *J. Exp. Med.* 178, 27-47; Sette, A. et al (1989) *J. Immunol* 143, 1265-1267.) This is due to the structure of the peptide binding groove in MHC class II molecules, which is open at both ends and allows for overhang of longer peptides outside of the critical binding core. In contrast, the peptide binding groove in MHC class I molecules is closed at both ends, strictly limiting the length of possible binding peptides. (Brown, J.H. et al (1993) *Nature* 364, 33-39).

Strong and long lasting immunity depends in part on CD4<sup>+</sup> helper T cell functions. Therefore the identification of Class II-restricted melanoma antigens will broaden the immunotherapeutic approaches to treating and/or prophylaxing against melanoma.

#### SUMMARY OF THE INVENTION

This invention relates, in general, to MHC Class II restricted melanoma antigens recognized by CD4<sup>+</sup>T-lymphocytes and the nucleic acid sequences encoding these antigens. This invention also provides therapeutic uses for the nucleic acid sequences, proteins or peptides described herein. In addition, this invention provides a method for identifying additional Class II restricted melanoma antigens.

It is a general object of the present invention to provide proteins, polypeptides or peptides which encode for Class II restricted melanoma antigens.

- 4 -

° It is another object of this invention to provide a recombinant molecule comprising a vector and all or part of the nucleic acid sequence encoding for a Class II restricted melanoma antigen.

5 It is another object of this invention to produce recombinant proteins or peptides encoded by all or part of the nucleic acid sequence encoding for a Class II restricted melanoma antigen.

10 It is a further object of this invention to provide methods for prophylactic or therapeutic uses for the Class II restricted melanoma antigens.

It is also an object of this invention to provide melanoma vaccines comprising all or part of the Class II restricted melanoma antigens.

15 It is a further object of this invention to provide immunogenic peptides demonstrated to be Class II restricted melanoma antigens for use in vaccines.

It is a particular object of this invention to provide tyrosinase peptides which are Class II restricted melanoma antigens.

20 In addition, it is another object of this invention to provide multivalent vaccines comprising at least one Class II restricted melanoma antigen and at least one other immunogenic molecule capable of eliciting an immune response in a mammal to melanoma antigens.

25 It is another object of this invention to provide a method for preventing or treating melanoma utilizing Class II restricted melanoma antigens in gene therapy protocols.

30 It is a further object of this invention to provide peptides derived from a tyrosinase protein sequence for use in vaccines.

35 It is yet another object of this invention to provide a method of prophylactic or therapeutic immunization for melanoma using the vaccines described herein.

- 5 -

It is a further object of this invention to provide a method of identifying Class II restricted melanoma antigens that would constitute potential targets for immunotherapy.

#### DESCRIPTION OF THE FIGURES

5 Figure 1 shows that the response of CD4<sup>+</sup> tumor infiltrating lymphocytes (TIL) from patient number 1088 to autologous melanoma cells is HLA-DR restricted. TIL ( $1 \times 10^6$ /ml) were cultured in the presence of autologous Epstein Barr Virus (EBV)-B cells ( $1 \times 10^6$ /ml) alone, or B  
10 cells pulsed with a lysate of autologous cultured melanoma cells ( $7 \times 10^5$  cell equivalents/ml). Secretion of GM-CSF (granulocyte/macrophage-colony-stimulating factor) following autologous tumor stimulation was significantly inhibited by the monoclonal antibody (mAb) L243 (anti-HLA-  
15 DR) and IVA12 (anti-HLA-DR,-DP,-DQ).

Figure 2 shows CD4<sup>+</sup> TIL 1088 recognize lysates of autologous and allogeneic cultured melanoma lines presented by autologous EBV-B cells, indicating a commonly expressed melanoma antigen. TIL cultured for 58 days in  
20 the presence of IL-2 were incubated for 20 hours (h) with tumor-pulsed 1088-EBV. GM-CSF secretion was measured by ELISA. All cells were at  $1 \times 10^6$ /ml.

Figure 3 shows CD4<sup>+</sup> TIL 1088 react with melanoma cells and normal melanocytes, but not with tumors of other  
25 histologies. TIL secreted significant amounts of GM-CSF when stimulated with autologous EBV-B cells pulsed with lysates of the autologous 1088-mel or normal melanocytes FM 902 and FM 907, but not pulsed with lysates of colon cancers, breast cancers, or Ewing's sarcomas. All cells  
30 were  $10^6$  per ml. Ca, cancer; Sa; sarcoma

Figure 4 shows that CD4<sup>+</sup> TIL 1088 recognize two different tyrosinase peptides. HLA-DRB1\*0401 was identified as the presenting molecule for both Ty 56-70 and Ty 448-462. CD4<sup>+</sup>TIL 1088 recognition was assessed by  
35 GM-CSF secretion. Peptides HA 307-319 and MT(65) 3-13

- 6 -

° were used as inhibitors for B1\*0401 and B1\*0301, respectively.

Figure 5 shows the MHC Class II restriction of tyrosinase peptide recognition by CD4<sup>+</sup> TIL 1088 cells. The antigen presenting cells were pulsed with autologous melanoma lysate or 100  $\mu$ M peptide overnight then washed  
5 prior to assay, all cells at  $1 \times 10^6$ /ml. HLA-DR-B1\*0401 was identified as the presenting MHC molecule for Ty 56-70 and Ty 448-462.

Figure 6 shows in three separate experiments, CD4<sup>+</sup>T cells from patient 1088 showed specific recognition  
10 of the Ty 56-70 peptide. Mutated and truncated peptides were used to identify the primary anchor (binding) residues within Ty 56-70. Recognition was measured as GM-CSF secretion by T cells cocultured for 24 hours with  
15 peptide-pulsed autologous EBV-transformed B cells.

Figure 7 shows the tyrosinase (Ty) peptides Ty 56-70 and Ty 448-462 with the P1 and P6 anchor positions boxed. The number designation of the peptide (e.g. Ty 56-70) indicates the amino acids spanned by the peptide, with  
20 1 being the methionine encoded by the initiation codon.

Figure 8 shows CD4<sup>+</sup>1088 TIL recognize modified tyrosinase 56-70 (Ty 56-70) peptides better than the nonmutated or non-modified peptide. (Background TIL + EBV = 118 pg/ml, subtracted.)

25 Figure 9 shows in two experiments, Ty 448-462 was specifically recognized by CD4<sup>+</sup>T cells from patient 1088. Mutated and truncated peptides were used to identify Y451 as the primary P1 binding residue within Ty 448-462.

30 Figure 10 shows CD4<sup>+</sup> TIL 1088 recognize truncated tyrosinase 448-462 peptides, Ty 449-462 and Ty 450-462, better than Ty 448-462. (Background TIL + EBV = 196 pg/ml, subtracted).

35 Figure 11 shows the P6 anchor position for Ty 448-462 is D456. This was presumed, based on identifying



- 7 -

Y451 as the P1 anchor. D is not an optimal residue in this position, and a valine substitution led to markedly enhanced CD4<sup>+</sup> T cell recognition. When two favorable modifications of Ty 448-462 were combined in a single modified peptide (Ty 450-462, D456 → V), recognition was enhanced even more (see Figure 12).

Figure 12 shows CD4<sup>+</sup> TIL 1088 recognize modified Tyrosinase 448-462 peptides better than the non-modified or nonmutated peptide (Background TIL + EBV = 99 pg/ml, subtracted).

#### DETAILED DESCRIPTION OF THE INVENTION

Major Histocompatibility Complex (MHC) is a generic designation meant to encompass the histocompatibility antigen systems described in different species, including the human leucocyte antigens (HLA).

The term melanoma includes, but is not limited to, melanomas, metastatic melanomas, melanomas derived from either melanocytes or melanocyte related nevus cells, melanocarcinomas, melanoepitheliomas, melanomasarcomas, melanoma in situ, superficial spreading melanoma, nodular melanoma, lentigo maligna melanoma, acral lentiginous melanoma, invasive melanoma or familial atypical mole and melanoma (FAM-M) syndrome. Such melanomas in mammals may be caused by, chromosomal abnormalities, degenerative growth and developmental disorders, mitogenic agents, ultraviolet radiation (UV), viral infections, inappropriate tissue expression of a gene, alterations in expression of a gene, and presentation on a cell, or carcinogenic agents. The aforementioned melanomas can be diagnosed, assessed or treated by methods described in the present application.

Immunogenic peptide includes, but is not limited to, an antigenic peptide capable of causing or stimulating a cellular or humoral immune response. Such peptides may also be reactive with antibodies.

- 8 -

This invention provides MHC-Class II restricted melanoma antigens. Such antigens may be the complete protein encoded by a gene, or portions thereof or polypeptides or peptides derived from a protein sequence. Such antigens may be expressed in normal or disease tissues. By way of example, Class II restricted melanoma antigens may be derived from the tyrosinase amino acid sequences. Examples of immunogenic tyrosinase sequences that may be used include, but are not limited to, GenBank accession numbers J03581, U01873, Y00819, and M27160, (Kwon, et al., (1987) *PNAS* 84:7473-7477; Brichard, V. et al., (1993) *J. Exp. Med.* 178:489-495; Bouchard, B. et al. (1989) *J. Exp. Med.* 169:2029-2042; and Shibihara, S. et al., (1988) *J. Exp. Med.* 156:403-414; all herein incorporated by reference). The Class II restricted melanoma antigen may comprise the entire tyrosinase sequence or portions thereof. Examples of immunogenic tyrosinase peptides recognized by CD4<sup>+</sup> T cells include, but are not limited to, QNILLSNAPLG PQFP (Ty 56-70) (SEQ ID NO: 1), NILLSNAPLG PQFP (Ty 57-70) (SEQ ID NO: 2), DYSYLQSDPDSFQD (Ty 448-462) (SEQ ID NO: 6), YSYLQSDPDSFQD (Ty 449-462) (SEQ ID NO: 13), and SYLQSDPDSFQD (Ty 450-462) (SEQ ID NO: 14) (Peptides are presented in single letter code). Also intended to be encompassed by this invention are proteins or polypeptides comprising these immunogenic peptide sequences. Persons of ordinary skill in the art will recognize that these peptides could be shortened to a minimal MHC Class II binding core of 9 or 10 amino acids by truncating the amino and/or carboxy termini of these peptides or one could lengthen these peptides by adding flanking sequences at either the carboxy or amino terminus of the peptides or at both termini of the peptide. By way of example, such a peptide may range in size from about 9 amino acids to about 34 amino acids.

35

- 9 -

° This invention further includes analogs of these immunogenic peptides derived from the tyrosinase amino acid sequence. The term analog includes any peptide which displays the functional aspects of these immunogenic peptides. The term analog also includes conservative substitution or chemical derivative of the peptides as described above. These peptides may be synthetically or recombina<sup>5</sup>ntly produced by conventional methodology.

The term "analog" includes any polypeptide having an amino acid residue sequence substantially identical to the sequences described herein in which one or more residues have been conservatively substituted with a functionally similar residue and which displays the functional aspects of the peptides as described herein. Examples of conservative substitutions include the substitution of one non-polar (hydrophobic) residue such as isoleucine, valine, leucine or methionine for another, the substitution of one polar (hydrophilic) residue for another such as between arginine and lysine, between glutamine and asparagine, between glycine and serine, the substitution of one basic residue such as lysine, arginine or histidine for another, or the substitution of one acidic residue, such as aspartic acid or glutamic acid for another. <sup>15</sup>

The phrase "conservative substitution" also includes the use of a chemically derivatized residue in place of a non-derivatized residue. "Chemical derivative" refers to a subject polypeptide having one or more residues chemically derivatized by reaction of a functional side group. Examples of such derivatized molecules include for example, those molecules in which free amino groups have been derivatized to form amine hydrochlorides, p-toluene sulfonyl groups, carbobenzoxy groups, t-butyloxycarbonyl groups, chloroacetyl groups or formyl groups. Free carboxyl groups may be derivatized to form salts, methyl and ethyl esters or other types of <sup>35</sup>

- 10 -

° esters or hydrazides. Free hydroxyl groups may be derivatized to form O-acyl or O-alkyl derivatives. The imidazole nitrogen of histidine may be derivatized to form N-im-benzylhistidine. Also included as chemical derivatives are those proteins or peptides which contain one or more naturally-occurring amino acid derivatives of the twenty standard amino acids. For examples: 4-hydroxyproline may be substituted for proline; 5-hydroxylysine may be substituted for lysine; 3-methylhistidine may be substituted for histidine; homoserine may be substituted for serine; and ornithine may be substituted for lysine. Proteins, polypeptides or proteins of the present invention also include any polypeptide having one or more additions and/or deletions or residues relative to the sequence of a polypeptide or peptide whose sequence are described herein, so long as the requisite activity is maintained.

In yet another embodiment of this invention, Class II restricted peptides derived from a tyrosinase sequence are modified to increase immunogenicity by enhancing the binding of the peptide to the MHC Class II molecule with which the peptide is associated when presented to CD4<sup>+</sup> T cells, or by enhancing binding of the peptide to the T cell receptor of the CD4<sup>+</sup> T cells. By way of example, modifications may include the substitution, deletion or addition, of one or more amino acids within the peptide sequence, or insertion of amino acids within the given peptide sequence or derivitization of existing amino acids within the given peptide sequence or mutation of the amino acids within the given peptide sequence. Examples of modified QNILLSNAPLGPQFP (Ty 56-70) (SEQ ID NO: 1) peptides include, but are not limited to, QNILLSNAPVGPQFP (L65→V) (SEQ ID NO: 3), QNILLSNVPGPQFP (A63→V and L65→V) (SEQ ID NO: 4), and QNILLSNVPLGPQFP (A63→V) (SEQ ID NO: 5) (See Figure 6). Examples of modified DYSYLQSDPDSFQD (Ty 448-462) (SEQ ID NO: 6)

- 11 -

° peptides include, but are not limited to DYSYLQSDPDSSQD (F460→S) (SEQ ID NO: 7), DQSYLQSDPDSPSFQD (Y449→Q) (SEQ ID NO: 8), DFSYLQSDPDSPSFQD (Y449→F) (SEQ ID NO: 9), DYSFLQSDPDSPSFQD (Y451→F) (SEQ ID NO: 10), DYSYLQDSVPDSFQD (D456→V) (SEQ ID NO: 11), and SYLQDSVPDSFQD (Ty450-462, 5 D456→V) (SEQ ID NO: 12) (see Figures 9 and 11).

Preferably the modifications are performed within the Class II core binding of the tyrosinase peptides. In a preferred modification at least one amino acid is substituted or replaced in the given binding core of the immunogenic peptide sequence. Any amino acid 10 composing the given binding core of the immunogenic peptide sequence may be modified in accordance with this invention. Any amino acid may be used to substitute or replace a given amino acid within the binding core of the immunogenic peptide sequence. Modification may occur at 15 any amino acid position within the binding core of an immunogenic tyrosinase peptide. Modified peptides is intended to include any modified peptide exhibiting enhanced binding with the MHC Class II molecule with which it is associated when presented to the CD4<sup>+</sup> T cell. Also 20 intended to be encompassed by this invention are proteins or polypeptides comprising or including these peptide sequences. By way of example such proteins or polypeptides may have additional sequences such as 25 flanking sequences either at the carboxy or amino terminus of the peptide or both.

By way of example, the Class II restricted tyrosinase antigens may be recognized by CD4<sup>+</sup>T cells in the context of HLA-DR, in particular HLA-DRB1\*0401. The 30 core binding sequence of a Class II restricted antigen is about 9 amino acids in length. Preferably for enhanced binding of the peptide to HLA-DRB1\*0401 the first position in the core amino acid sequence is an aromatic or aliphatic hydrophobic amino acid. The sixth position may 35 be any hydrophobic amino acid such as, but not limited to,

- 12 -

leucine, isoleucine, valine, methionine, or a hydroxyl amino acid, such as serine or threonine (Sette, A. et al (1993) *J. Immunol.* 151, 3163-3170; Rammensee, H.G. et al (1995) *Immunogenetics* 41:178-228, both herein incorporated by reference).

The fourth, seventh and ninth positions of the 9 amino acid binding core sequence of the immunogenic peptide may also be substituted or replaced. Examples of amino acids that may be used at the fourth position of the peptide include, but are not limited to, any hydrophobic amino acid or aspartic or glutamic acid. The seventh position may be any polar, charged or aliphatic amino acid. Examples of amino acids that may be used include but are not limited to aspartic acid, alanine, serine, valine, histidine, proline, asparagine, methionine, threonine, leucine and isoleucine. The ninth position of the peptide may be any polar or aliphatic amino acid. Examples of such amino acids include but are not limited to alanine, serine, glutamine, glycine, leucine, valine, and threonine.

Examples of Class II restricted tyrosinase peptides whose core sequence may be modified in accordance with the present embodiment include, but is not limited to QNILLSNAPLGPQFP (Ty 56-70), NILLSNAPLGPQFP (Ty 57-70), DYSYLQSDPDSPDFQD (Ty 448-462), YSYLQSDPDSPDFQD (Ty 449-462), and SYLQSDPDSPDFQD (Ty 450-462). Examples of modified tyrosinase peptides whose core binding sequence may further be modified include, but is not limited to, QNILLSNAPVGPQFP (Ty56-70, L65→V), QNILLSNVPVGPQFP (Ty 56-70, A63→V and L65→V), QNILLSNVPLGPQFP (Ty 56-70, A63→V), DYSYLQSDPDSSQD (Ty 448-462, F460→S), DQSYLQSDPDSPDFQD (Ty 448-462, Y449→Q), DFSYLQSDPDSPDFQD (Ty 448-462, Y449→F), DYSFLQSDPDSPDFQD (Ty 448-462, Y451→F), DYSYLQSDVPDSPDFQD (Ty 448-462, D456→V), and SYLQSDVPDSPDFQD (Ty450-462, D456→V).

35

- 13 -

By way of example modified Class II-restricted tyrosinase peptides derived from the tyrosinase sequence may have a binding core sequence according to the formula  $X_1LLX_2NX_3X_4LX_5$ , (SEQ ID NO: 15), or  $X_1LQX_2SX_3X_4DX_5$ , (SEQ ID NO: 16), wherein:

$X_1$  may be any hydrophobic amino acid, either aromatic or aliphatic. Examples of amino acids that may be used include, but are not limited to, leucine, isoleucine, methionine, valine, tryptophan, phenylalanine, or tyrosine. The  $X_1$  position corresponds to Ty 58 in peptide Ty56-70, or Ty 451 in peptide Ty 448-462.

$X_2$  may be any hydrophobic amino acid, or aspartic acid or glutamic acid. Examples of amino acids that may be used include, but are not limited to, phenylalanine, tryptophan, leucine, isoleucine, valine, alanine, aspartic acid or glutamic acid.

$X_3$  may be any hydrophobic amino acid, or hydroxyl amino acids. Examples of amino acids that may be used include, but are not limited to, leucine, isoleucine, methionine, valine, serine or threonine.

$X_4$  may be any polar, charged or aliphatic amino acid. Examples of amino acids that may be used include, but are not limited to, aspartic acid, alanine, serine, valine, histidine, proline, asparagine, methionine, threonine, leucine, and isoleucine.

$X_5$  may be any polar, or aliphatic amino acid. By way of example amino acids that may be used on this position include, but are not limited to, alanine, serine, glutamine, glycine, leucine, valine, and threonine.

This invention further includes analogs of these modified peptides derived from the tyrosinase sequence. The term analog is intended to include any peptide which displays the functional aspects of these modified peptides as described above. These modified peptides may be synthetically or recombinantly produced by conventional methods. Also intended to be encompassed by this

- 14 -

invention are proteins or polypeptides including these peptide sequences. By way of example such proteins or polypeptides may have additional flanking sequences at either the carboxy or amino terminus of the peptide or at both termini or may be truncated to a minimal 9 amino acid MHC Class II binding core.

This invention also relates to nucleic acid encoding the Class II restricted immunogenic tyrosinase peptides or modified peptides of this invention. It is, however, understood by one skilled in the art that due to the degeneracy of the genetic code variations in nucleic acid sequences will still result in a DNA sequence capable of encoding antigens described herein. Such DNA sequences are therefore functionally equivalent to the sequences intended to be encompassed by the invention. Allelic variations in a given species of the nucleic acid sequence encompassed by this invention are also intended to be encompassed by the present invention. By way of example nucleic acid sequences encoding the tyrosinase or modified tyrosinase peptides QNILLSNAPLGPQFP (Ty 56-70), NILLSNAPLGPQFP (Ty 57-70), DYSYLQSDPDSFQD (Ty 448-462), YSYLQSDPDSFQD (Ty 449-462), SYLQSDPDSFQD (Ty 450-462), QNILLSNAPVGPQFP (Ty 56-70, L65→V), QNILLSNVPGPQFP (Ty 56-70, A63→V and L65→V), QNILLSNVPLGPQFP (Ty 56-70, A63→V) DYSYLQSDPDSSQD (Ty 448-462, F460→S), DQSYLQSDPDSFQD (Ty 448-462, Y449→Q), DFSYLQSDPDSFQD (Ty 448-462, Y449→F), DYSFLQSDPDSFQD (Ty 448-462, Y451→F), DYSYLQDSVPDSFQD (Ty-448-462, D456→V), and SYLQDSVPDSFQD (Ty450-462, D456→V) or analogs thereof are intended to be encompassed by this invention.

This invention also provides a recombinant DNA molecule comprising all or part of the nucleic acid sequence encoding a Class II melanoma antigen and a vector. Expression vectors suitable for use in the present invention comprise at least one expression control element operationally linked to the nucleic acid sequence.



- 15 -

° The expression control elements are inserted in the vector to control and regulate the expression of the nucleic acid sequence. Examples of expression control elements include, but are not limited to, lac system, operator and promoter regions of phage lambda, yeast promoters and  
5 promoters derived from polyoma, adenovirus, retrovirus or SV40. Additional preferred or required operational elements include, but are not limited to, leader sequence, termination codons, polyadenylation signals and any other sequences necessary or preferred for the appropriate  
10 transcription and subsequent translation of the nucleic acid sequence in the host system. It will be understood by one skilled in the art that the correct combination of required or preferred expression control elements will depend on the host system chosen. It will further be  
15 understood that the expression vector should contain additional elements necessary for the transfer and subsequent replication of the expression vector containing the nucleic acid sequence in the host system. Examples of such elements include, but are not limited to, origins of  
20 replication and selectable markers. It will further be understood by one skilled in the art that such vectors are easily constructed using conventional methods (Ausubel et al., (1987) in "Current Protocols in Molecular Biology", John Wiley and Sons, New York, New York) or are  
25 commercially available.

Another aspect of this invention relates to a host organism into which a recombinant expression vector containing all or part of the nucleic acid sequence encoding for a Class II melanoma antigens has been  
30 inserted. The host cells transformed with the nucleic acid sequences encompassed by this invention include eukaryotes, such as animal, plant, insect and yeast cells and prokaryotes, such as E. coli. The means by which the vector carrying the gene may be introduced into the cell  
35 include, but are not limited to, microinjection,

- 16 -

° electroporation, transduction, or transfection using DEAE-dextran, lipofection, calcium phosphate or other procedures known to one skilled in the art (Sambrook et al. (1989) in "Molecular Cloning. A Laboratory Manual", Cold Spring Harbor Press, Plainview, New York).

5 In a preferred embodiment, eukaryotic expression vectors that function in eukaryotic cells are used. Examples of such vectors include, but are not limited to, retroviral vectors, vaccinia virus vectors, adenovirus vectors, herpes virus vector, fowl pox virus vector, 10 plasmids, such as pCDNA3 (Invitrogen, San Diego, CA) or the baculovirus transfer vectors. Preferred eukaryotic cell lines include, but are not limited to, COS cells, CHO cells, HeLa cells, NIH/3T3 cells, 293 cells (ATCC# CRL1573), T2 cells, dendritic cells, monocytes or Epstein- 15 Barr Virus transformed B cells. In a preferred embodiment the recombinant protein expression vector is introduced into mammalian cells, such as NIH/3T3, COS-7, CHO, 293 cells (ATCC #CRL 1573), T2 cells, dendritic cells, or monocytes to ensure proper processing and modification of 20 the protein.

The recombinant protein expressed by the host cells can be obtained as a crude lysate or can be purified by standard protein purification procedures known in the art which may include differential precipitation, 25 molecular sieve chromatography, ion-exchange chromatography, isoelectric focusing, gel electrophoresis, affinity, and immunoaffinity chromatography and the like. (Ausubel et. al., (1987) in "Current Protocols in Molecular Biology" John Wiley and Sons, New York, New 30 York). In the case of immunoaffinity chromatography, the recombinant protein may be purified by passage through a column containing a resin which has bound thereto antibodies specific for the tyrosinase protein (Ausubel et. al., (1987) in "Current Protocols in Molecular 35 Biology" John Wiley and Sons, New York, New York).

- 17 -

This invention further includes an antibody or antibodies reactive with the Class II restricted melanoma antigens described. The antibodies may be monoclonal and polyclonal and are made by conventional methods known to those skilled in the art. In addition, the protein or nucleic acid sequences of the Class II restricted melanoma antigens described herein, may be used diagnostically to screen for the presence, absence or alteration in expression of these antigens using immunoassays or nucleic acid probes.

The Class II restricted melanoma antigens of this invention, or analogs thereof may be used as a vaccine either prophylactically or therapeutically. When provided prophylactically the vaccine is provided in advance of any evidence of melanoma. The prophylactic administration of the Class II restricted melanoma antigen vaccine should serve to prevent or attenuate melanoma in a mammal. In a preferred embodiment mammals, preferably human, at high risk for melanoma are prophylactically treated with the vaccines of this invention. Examples of such mammals include, but are not limited to, humans with a family history of melanoma, humans with a history of atypical moles, humans with a history of FAM-M syndrome or humans afflicted with melanoma previously resected and therefore at risk for reoccurrence. When provided therapeutically, the vaccine is provided to enhance the patient's own immune response to the tumor antigen present on the melanoma or metastatic melanoma. The vaccine, which acts as an immunogen, may be a cell, cell lysate from cells transfected with a recombinant expression vector, cell lysates from cells transfected with a recombinant expression vector encoding for the Class II restricted melanoma antigen, or a culture supernatant containing the expressed protein. Alternatively, the immunogen is a partially or substantially purified recombinant protein, peptide or analog thereof encoding for a Class II

- 18 -

restricted melanoma antigen. The proteins or peptides may be conjugated with lipoprotein or administered in liposomal form or with adjuvant using conventional methodologies. Examples of Class II restricted tyrosinase peptides or modified tyrosinase peptides that may be used include, but are not limited to, QNILLSNAPLGPQFP (Ty 56-70), NILLSNAPLGPQFP (Ty 57-70), DYSYLQSDPDSFQD (Ty 448-462), YSYLQSDPDSFQD (Ty 449-462), SYLQSDPDSFQD (Ty 450-462), QNILLSNAPVGPQFP (Ty 56-70, L65→V), QNILLSNVPGPQFP (Ty 56-70, A63→V and L65→V), QNILLSNVPLGPQFP (Ty 56-70, A63→V), DYSYLQSDPDSFQD (Ty 448-462, F460→S), DQSYLQSDPDSFQD (Ty 448-462, Y449→Q), DFSYLQSDPDSFQD (Ty 448-462, Y449→F), DYSFLQSDPDSFQD (Ty 448-462, Y451→F), DYSYLQSDVPDSFQD (Ty 448-462, D456→V), and SYLQSDVPDSFQD (Ty 450-462, D456→V) or analogs thereof. The tyrosinase protein or tyrosinase peptides having the modified binding core sequences described herein may also be used.

While it is possible for the immunogen to be administered in a pure or substantially pure form, it is preferable to present it as a pharmaceutical composition, formulation or preparation.

The formulations of the present invention, both for veterinary and for human use, comprise an immunogen as described above, together with one or more pharmaceutically acceptable carriers and, optionally, other therapeutic ingredients. The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof. The formulations may conveniently be presented in unit dosage form and may be prepared by any method well-known in the pharmaceutical art.

All methods include the step of bringing into association the active ingredient with the carrier which constitutes one or more accessory ingredients. In general, the formulations are prepared by uniformly and

- 19 -

- ° intimately bringing into association the active ingredient with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product into the desired formulation.

Formulations suitable for intravenous,  
5 intramuscular, subcutaneous, or intraperitoneal administration conveniently comprise sterile aqueous solutions of the active ingredient with solutions which are preferably isotonic with the blood of the recipient. Such formulations may be conveniently prepared by  
10 dissolving solid active ingredient in water containing physiologically compatible substances such as sodium chloride (e.g. 0.1-2.0M), glycine, and the like, and having a buffered pH compatible with physiological conditions to produce an aqueous solution, and rendering  
15 said solution sterile. These may be present in unit or multi-dose containers, for example, sealed ampoules or vials.

The formulations of the present invention may incorporate a stabilizer. Illustrative stabilizers are  
20 polyethylene glycol, proteins, saccharides, amino acids, inorganic acids, and organic acids which may be used either on their own or as admixtures. These stabilizers are preferably incorporated in an amount of about 0.11 about 10,000 parts by weight per part by weight of  
25 immunogen. If two or more stabilizers are to be used, their total amount is preferably within the range specified above. These stabilizers are used in aqueous solutions at the appropriate concentration and pH. The specific osmotic pressure of such aqueous solutions is  
30 generally in the range of about 0.1 to about 3.0 osmoles, preferably in the range of about 0.8 to about 1.2. The pH of the aqueous solution is adjusted to be within the range of about 5.0 to about 9.0, preferably within the range of 6-8. In formulating the immunogen of the present  
35 invention, anti-adsorption agent may be used.

- 20 -

Additional pharmaceutical methods may be employed to control the duration of action. Controlled release preparations may be achieved through the use of polymer to complex or absorb the proteins or their derivatives. The controlled delivery may be exercised by selecting appropriate macromolecules (for example polyester, polyamino acids, polyvinyl, pyrrolidone, ethylenevinylacetate, methylcellulose, carboxymethylcellulose, or protamine sulfate) and the concentration of macromolecules as well as the methods of incorporation in order to control release. Another possible method to control the duration of action by controlled-release preparations is to incorporate the tyrosinase protein, peptides and analogs thereof into particles of a polymeric material such as polyesters, polyamino acids, hydrogels, poly(lactic acid) or ethylene vinylacetate copolymers. Alternatively, instead of incorporating these agents into polymeric particles, it is possible to entrap these materials in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxy-methylcellulose or gelatin-microcapsules and poly(methylmethacrylate) microcapsules, respectively, or in colloidal drug delivery systems, for example, liposomes, albumin microspheres, microemulsions, nanoparticles, and nanocapsules or in macroemulsions.

When oral preparations are desired, the compositions may be combined with typical carriers, such as lactose, sucrose, starch, talc magnesium stearate, crystalline cellulose, methyl cellulose, carboxymethyl cellulose, glycerin, sodium alginate or gum arabic among others.

The proteins of the present invention may be supplied in the form of a kit, alone, or in the form of a pharmaceutical composition as described above.

35

- 21 -

°  
Vaccination can be conducted by conventional methods. For example, the immunogen can be used in a suitable diluent such as saline or water, or complete or incomplete adjuvants. Further, the immunogen may or may not be bound to a carrier to make the protein immunogenic or enhance the protein's immunogenicity. Examples of such carrier molecules include but are not limited to bovine serum albumin (BSA), keyhole limpet hemocyanin (KLH), tetanus toxoid, and the like. The immunogen also may be coupled with lipoproteins or administered in liposomal form or with adjuvants. The immunogen can be administered by any route appropriate for antibody production such as intravenous, intraperitoneal, intramuscular, subcutaneous, and the like. The immunogen may be administered once or at periodic intervals until a significant titer of CD4<sup>+</sup> or CD8<sup>+</sup> T cell or antibodies directed against the Class II restricted melanoma antigen is obtained. The presence of cells may be assessed by measuring cytokine secretion in response to antigen-presenting cells pulsed with the immunogen. The antibody may be detected in the serum using conventional immunoassays.

The administration of the vaccine or immunogen of the present invention may be for either a prophylactic or therapeutic purpose. When provided prophylactically, the immunogen is provided in advance of any evidence or in advance of any symptom due to melanoma, or in patients rendered free of disease by conventional therapies but at significant risk for recurrence. The prophylactic administration of the immunogen serves to prevent or attenuate melanoma in a mammal. When provided therapeutically, the immunogen is provided at (or after) the onset of the disease or at the onset of any symptom of the disease. The therapeutic administration of the immunogen serves to attenuate the disease.

By way of example, a vaccine prepared using recombinant expression vectors may be used. To provide a

- 22 -

° vaccine to an individual a genetic sequence which encodes  
for all or part of the Class II restricted melanoma  
antigen is inserted into an expression vector, as  
described above, and introduced into the mammal to be  
immunized. Examples of vectors that may be used in the  
5 aforementioned vaccines include, but are not limited to,  
defective retroviral vectors, adenoviral vectors, vaccinia  
viral vectors, fowl pox viral vectors, or other viral  
vectors (Mulligan, R.C., (1993) Science 260:926-932). The  
viral vectors carrying the nucleic sequence can be  
10 introduced into a mammal either prior to any evidence of  
melanoma or to mediate regression of the disease in a  
mammal afflicted with melanoma. Examples of methods for  
administering the viral vector into the mammals include,  
but are not limited to, exposure of cells to the virus ex  
15 vivo, or injection of the retrovirus or a producer cell  
line of the virus into the affected tissue or intravenous  
administration of the virus. Alternatively the viral  
vector carrying all or part of the tyrosinase nucleic acid  
sequence encoding the Class II restricted melanoma antigen  
20 may be administered locally by direct injection into the  
melanoma lesion or topical application in a  
pharmaceutically acceptable carrier. Examples of nucleic  
acid sequences that may be used include, but are not  
limited to, nucleic acid sequence encoding the Class II  
25 tyrosinase restricted peptides or modified peptides  
QNILLSNAPLGPQFP (Ty 56-70), NILLSNAPLGPQFP (Ty 57-70),  
DYSYLQSDPDSFQD (Ty 448-462), YSYLQSDPDSFQD (Ty 449-462),  
SYLQSDPDSFQD (Ty 450-462), QNILLSNAPVGPQFP (Ty 56-70,  
L65→V), QNILLSNVPGPQFP (Ty 56-70, A63→V and L65→V),  
30 QNILLSNVPLGPQFP (Ty 56-70, A63→V), DYSYLQSDPDSSQD (Ty  
448-462, F460→S), DQSYLQSDPDSFQD (Ty 448-462, Y449→Q),  
DFSYLQSDPDSFQD (Ty 448-462, Y449→F), DYSFLQSDPDSFQD (Ty  
448-462, Y451→F), DYSYLQSDVPDSFQD (Ty 448-462, D456→V),  
and SYLQSDVPDSFQD (Ty450-462, D456-V) or analogs thereof.  
35 In addition, nucleic acid sequences encoding tyrosinase



- 23 -

peptides comprising the modified core binding sequences described herein may also be incorporated into recombinant vectors. The quantity of viral vector, carrying the nucleic acid sequence encoding for the Class II restricted melanoma antigen, to be administered is based on the titer of virus particles. By way of example, a range of the immunogen to be administered may be about  $10^6$  to about  $10^{11}$  virus particles per mammal, preferably a human. After immunization the efficacy of the vaccine can be assessed by production of antibodies or immune cells that recognize the antigen, as assessed by specific cytokine production or by tumor regression. One skilled in the art would know the conventional methods to assess the aforementioned parameters. If the mammal to be immunized is already afflicted with melanoma or metastatic melanoma the vaccine can be administered in conjunction with other therapeutic treatments. Examples of other therapeutic treatment includes, but are not limited to, adoptive T cell immunotherapy, coadministration of cytokines or other therapeutic drugs for melanoma.

Alternatively all or parts thereof of a substantially or partially purified tyrosinase protein corresponding to the Class II restricted melanoma antigen or polypeptides or peptides may be administered as a vaccine in a pharmaceutically acceptable carrier. By way of example, ranges of protein polypeptides or peptides to be administered may be 0.001 to about 100 mg per patient, preferred doses are about 0.01 to about 10mg per patient. In a preferred embodiment, tyrosinase Class II restricted peptide melanoma antigens or analogs thereof or modified tyrosinase peptides are administered therapeutically or prophylactically to a mammal in need of such treatment. By way of example, doses may be about 0.001 mg to about 100 mg, preferred doses are about 0.01 mg to about 10mg. The peptide may be synthetically or recombinantly produced. Immunization may be repeated as necessary,

- 24 -

until a sufficient titer of anti-immunogen antibody or reactive CD4<sup>+</sup> or CD8<sup>+</sup> T cells has been obtained.

In yet another alternative embodiment a viral vector, such as a retroviral vector, can be introduced into mammalian cells. Examples of mammalian cells into which the retroviral vector can be introduced include, but are not limited to, primary mammalian cultures or continuous mammalian cultures, COS cells, NIH3T3, or 293 cells (ATTC #CRL 1573), B cell, dendritic or monocytic cell cultures. The means by which the vector carrying the gene may be introduced into a cell includes, but is not limited to, microinjection, electroporation, transfection or transfection using DEAE dextran, lipofection, calcium phosphate or other procedures known to one skilled in the art (Sambrook et al. (eds) (1989) in "Molecular Cloning. A Laboratory Manual", Cold Spring Harbor Press, Plainview, New York). The mammalian cells expressing the Class II restricted melanoma antigen can be administered to mammals and serve as a vaccine or immunogen. Examples of how the cells expressing Class II restricted melanoma antigens can be administered include, but is not limited to, subcutaneous, intravenous, intraperitoneal or intralesional. In a preferred embodiment, the nucleic acid sequence corresponding to Class II restricted peptides or modified tyrosinase peptides is inserted into expression vector and introduced into the mammalian cells. By way of example, the peptides that may be used include, but are not limited to, QNILLSNAPLGPQFP (Ty 56-70), NILLSNAPLGPQFP (Ty 57-70), DYSYLQSDPDSFQD (Ty 448-462), YSYLQSDPDSFQD (Ty 449-462), SYLQSDPDSFQD (Ty 450-462), QNILLSNAPVGPQFP (Ty 56-70, L65→V), QNILLSNVPGPQFP (Ty 56-70, A63→V and L65→V), QNILLSNVPLGPQFP (Ty 56-70, A63→V), DYSYLQSDPDSSQD (F460→S), DQSYLQSDPDSFQD (Ty 448-462, Y449→Q), DFSYLQSDPDSFQD (Ty 448-462, Y449→F), DYSFLQSDPDSFQD (Ty 448-462, Y451→F), DYSYLQSDVPDSFQD (Ty 448-462, D456→V), and SYLQSDVPDSFQD (Ty450-462, D456-V) or

- 25 -

° analogs thereof. Nucleic acid sequences encoding tyrosinase peptides having the binding core sequences provided may also be used. Conventional methods would be used to evaluate the immune response of the patient to determine the efficiency of the vaccine.

5 In yet another embodiment of this invention, the Class II restricted tyrosinase protein, peptides or modified peptides or analogs thereof may be exposed to dendritic cells cultured in vitro. The cultured dendritic cells provide a means of producing CD4<sup>+</sup> T cell dependent  
10 antigens comprised of dendritic cell modified antigen or dendritic cells pulsed with antigen, in which the protein antigen is processed and expressed on the antigen activated dendritic cell. The antigen activated dendritic cells or processed dendritic cell antigens may be used as  
15 immunogens for vaccines or for the treatment of melanoma. Alternatively the dendritic cells may present peptide antigens which have been pulsed on externally. The dendritic cells should be exposed to antigen for sufficient time to allow the antigens to bind directly to  
20 their surface MHC Class II molecules, or to be internalized and presented on the dendritic cells surface. The resulting dendritic cells or the dendritic cell processed antigens can than be administered to an individual in need of therapy. Such methods are described  
25 in Steinman et al. (WO93/208185) and in Banchereau et al. (EPO Application 0563485A1) which are incorporated herein by reference. Monocytes, B cells, or Langerhans cells may be substituted for dendritic cells.

In yet another embodiment of this invention CD4<sup>+</sup>  
30 T cells isolated from individuals can be exposed to the Class II restricted melanoma antigen in vitro and then administered to a patient in need of such treatment in a therapeutically effective amount. Examples of where CD4<sup>+</sup>T-lymphocytes can be isolated, include but are not  
35 limited to, peripheral blood cells lymphocytes (PBL),

- 26 -

° lymph nodes, or tumor infiltrating lymphocytes (TIL). Such lymphocytes can be isolated from the individual to be treated or from a donor by methods known in the art and cultured in vitro (Kawakami, Y. et al. (1989) J. Immunol. 142: 2453-3461). The CD4<sup>+</sup>T-lymphocytes are cultured by  
5 methods known in the art. The lymphocytes are exposed to peptide or protein antigen for part or all of the culture duration, in the presence of antigen presenting cells. In a preferred embodiment the CD4<sup>+</sup>lymphocytes are exposed to the Class II restricted tyrosinase peptides or any  
10 tyrosinase sequence having the core peptide sequences described herein. By way of example, a concentration of about 1 to about 200 micrograms(ug)/ml peptides per 10<sup>7</sup> cells for all or part of the duration of lymphocyte culture may be used. After being sensitized to the  
15 peptide the T-lymphocytes are administered to the mammal in need of such treatment. Examples of how these sensitized CD4<sup>+</sup> T cells can be administered to the mammal include but are not limited to, intravenously, intraperitoneally or intralesionally. Parameters that may  
20 be assessed to determine the efficacy of these sensitized T-lymphocytes include, but are not limited to, production of immune cells in the mammal being treated or tumor regression. Conventional methods are used to assess these parameters. Such treatment can be given in conjunction  
25 with cytokines or gene modified cells (Rosenberg, S.A. et al. (1992) Human Gene Therapy, 3: 75-90; Rosenberg, S.A. et al. (1992) Human Gene Therapy, 3: 57-73). By way of example the MHC Class II restricted melanoma antigen may be administered in conjunction with GM-CSF to enhance  
30 uptake by professional antigen presenting cells in vivo.  
In yet another alternative embodiment Class II restricted melanoma antigens may be linked with MHC Class II molecules and administered either prophylactically or therapeutically to mammals. By way of example the  
35 tyrosinase peptides or modified peptides described herein

- 27 -

may be coupled with an MHC Class II molecule. Such coupling may be covalent, chemical, or genetic. By way of example, tyrosinase peptides may be genetically linked to the Class II  $\beta$  chain, HLA-DRB1\*0401, by a flexible peptide linker which allows the peptide to lie in the binding groove for recognition by T cells (Kozono, H et al (1994) Nature 369, 151-154, herein incorporated by reference). According to the promiscuous and degenerate nature of peptide binding to MHC Class II molecules, other DRB1 chains may be used as well (Sinigaglia, F. et al (1995) J. Exp. Med. 181, 449-451). Such single-chain Class II-MHC-peptide constructs may be used as vaccines, or may be used to raise reactive CD4<sup>+</sup> T cells in vitro.

In yet another alternative embodiment, the aforementioned compositions can be used to prepare antibodies to the Class II restricted melanoma antigens. The antibodies can be used directly as anti-melanoma therapeutic agents or as diagnostic reagents. Further, the antibodies can be made even more compatible with the host system by generating "humanized" chimeric antibodies (Morrison J. 1985, Science 229:1202 and Oi, et al. Biotechniques (1986) 4:214). Such antibodies can be generated by conventional methods.

In yet another embodiment of this invention, multivalent vaccines against one or more melanoma antigens are provided. Such multivalent vaccines may comprise at least one of the Class II restricted melanoma antigens described herein, preferably the tyrosinase immunogenic peptides disclosed herein or combinations thereof, combined with other known melanoma antigens or peptides derived from these antigens. By way of example, MHC Class II restricted tyrosinase peptides may be combined in a vaccine with MHC Class I restricted peptides derived from tyrosinase or other known melanoma associated proteins to create a multivalent vaccine capable of stimulating helper and cytotoxic immune cells. Examples of known melanoma

- 28 -

° antigens include, but are not limited to, MART-1, gp100, gp75, MAGE-1 and MAGE-3 or immunogenic peptides derived from these proteins.

5 In another embodiment, a method is provided for identifying the presence of Class II restricted antigenic proteins or epitopes of proteins, or of identifying new proteins encoding Class II restricted tumor associated antigens, such as, but not limited to, melanoma antigens. By way of example, the genes or nucleic acid sequences encoding Class I restricted melanoma antigens can be  
10 screened for the presence of a Class II restricted antigenic portions or epitopes of the protein encoded by these genes. In this embodiment, the method may comprise the steps of: (a) exposing a candidate antigen to antigen presenting cells (APC) for a period of time sufficient to  
15 allow the APC to take up and process the antigen; (b) incubating the APC of step (a) with CD4<sup>+</sup>T-lymphocytes; and (c) screening for recognition of the APC by the CD4<sup>+</sup> T cells (see Example 1).

In step (a) the candidate antigen may be  
20 presented to the APC by either stably or transiently expressing the gene for the candidate antigen in a eukaryotic or prokaryotic expression system. The antigen may then be presented to the APC as crude lysates of the cells expressing the candidate antigen or as purified  
25 protein products from the candidate antigen expressing cells. Alternatively a plurality of peptides based on the candidate protein amino acid sequence or based on a truncated protein sequence derived from experiments with serial truncations of the candidate gene may be exposed or  
30 incubated with the antigen presenting cell. It is preferred that peptides of about 15 to 20 amino acids be used.

Examples of APC that may be used in step (a) include, but are not limited to, antigen presenting cells  
35 such as EBV transformed B cell lines (Topalian et al.

- 29 -

(1994) *Int. J. Cancer* 58:69-79), monocytes and dendritic cells. Examples of how to assess recognition by the CD4<sup>+</sup> T cells incubated with the APC in step (c) include, but are not limited to, <sup>51</sup>CR release cytotoxicity assays (Cerundolo, V. et al. (1990) *Nature* 345:449-452.), cytokine secretion assays such as  $\gamma$ -IFN, GM-CSF or TNF secretion. (Schwartzentruber, D. et al., (1991) *J. of Immunology* 146:3674-3681), or proliferation assays.

Examples of proteins that may be screened for Class II restricted melanoma antigens includes, but are not limited to MART-1 (Kawakami, et al. (1994) *Proc. Natl. Acad. Sci.* 91:3575-3579), p15 (Robbins, P.F. et al (1995) *J. Immunol.* 154:5944-5950), MAGE-1 (VanderBruggen, *Science* 254:1643-1647), gp100 (Kawakami, et al. (1994) 91:6458-6462), gp75 (Wang, R-F et al (1995) *J. Exp. Med.* 181:799-804), and MAGE-3 (Gaugler, et al. (1994), *J. Exp. Med.* 179:921-930); all herein incorporated by reference.)

Alternatively, this method can be used to clone and identify new genes having CD4 recognized tumor antigens. By way of example, a cell expressing an unidentified tumor antigen would be assessed for CD4<sup>+</sup> T cell recognition by pulsing lysates of that cell onto antigen presenting cells (EBV-cells, monocytes, dendritic cells, etc.), and measuring cytokine secretion by T cells during coincubation. A DNA library from the tumor or other stimulatory cell would be expressed in a prokaryotic or eukaryotic host cell and screened according to the methods outlined above.

Also intended to be encompassed by this invention are Class II restricted tumor associated antigens, such as melanoma antigens obtained by these methods.

Veterinary uses are also intended to be encompassed by the compositions and therapeutic applications described herein.

- 30 -

All books, articles, and patents referenced herein are incorporated by reference. The following examples illustrate various aspects of the invention and in no way intended to limit the scope thereof.

## Example I

Human CD4<sup>+</sup> Cells Recognize a Shared  
Melanoma Antigen Encoded by the Tyrosinase Gene

MATERIALS AND METHODSLymphocyte Cultures and Clones.

TIL were cultured from enzymatically digested single cell suspensions of solid metastatic melanoma lesions as previously described (Topalian, S.L., et al. (1987) *J. Immunol. Methods.* 102, 127-141), in the presence of recombinant interleukin-2 (rIL-2; 6000 IU/ml (Chiron Corporation, Emeryville, CA)), Lymphocyte cultures were passaged in bulk for 4 to 6 weeks, and then CD4<sup>+</sup> and CD8<sup>+</sup> TIL subsets were purified by positive selection on tissue culture flasks carrying covalently bound anti-CD4 or anti-CD8 monoclonal antibodies (mAb) (Applied Immune Sciences, Menlo, Park, CA) (Morecki, S., et al. (1990) *J. Biol. Resp. Modif.* 9, 463-474). Uncloned CD4<sup>+</sup> TIL were tested in bioassays after 45-70 days of culture. CD4<sup>+</sup> T cell clones were established from a 35-day bulk TIL 1088 culture which was 52% CD4<sup>+</sup>. CD4<sup>+</sup> T cells were selected and then cloned by limiting dilution in microtiter plates, in the presence of 600 IU/ml IL-2, pooled allogeneic PBL from 3 donor's (total 3 x 10<sup>4</sup> cells/well, 3000 rad; 1 rad = 0.01Gy), and autologous Epstein Barr Virus (EBV)-B cells (1088-EBV 1 x 10<sup>4</sup> cells/well, 10,000 rad) pulsed with a freeze/thaw lysate of autologous tumor (1088-mel). Clones were restimulated weekly with allogeneic peripheral blood lymphocytes (PBL), 1088-EBV, and whole 1088-mel cells (1 x 10<sup>3</sup> cells/well, 30,000 rad). Clones used for bioassays were grown from 0.3 or 1 cell/well dilutions, and tested after 58-155 days of culture.



- 31 -

° B Cell Lines.

EBV-transformed B cell lines were established from the PBL of melanoma patients using standard techniques (Current Protocols In Immunology, Coligan, J. E. et al. (eds) Wiley and Sons, N.Y., N.Y. 7.22.1-7.22.3 (1994) herein incorporated by reference). These lines were maintained as suspension cultures in RPMI + 10% FCS. Tumors and Normal Melanocytes.

Melanoma cultures were established from fresh or cryopreserved single cell suspensions of metastatic lesions and maintained as adherent monolayers in RPMI + 10% FCS, as described (Topalian, S.L., et al. (1989) *J. Immunol.* 142, 3714-3725). Tumor clones were established by limiting dilution in 96-well flat bottom plates (Costar).

The normal melanocyte cultures FM 707, FM 708, FM 902, FM 906 and FM 907, generated from neonatal foreskin, were a generous gift of Dr. Meenhard Herlyn (Wistar Institute, Philadelphia, PA; Herlyn, M., et al. (1985) *Cancer Res.* 45, 5670-5676). Cultures were maintained in Melanocyte Growth Medium (MGM, Clonetics, San Diego, CA), a basal serum-free medium supplemented with recombinant basic fibroblast growth factor (1 ng/ml), insulin (5 ug/ml), hydrocortisone (0.5 ug/ml), PMA (10 ng/ml), bovine pituitary extract, gentamicin (50 ug/ml), and amphotericin B (50 ng/ml). Cells were removed from this medium for several days prior to bioassays.

The cultured colon carcinoma CY13 was a gift of Dr. J. Yannelli (NIH, Bethesda, MD). Colon carcinomas WiDr, LoVo, SW480; breast carcinomas ZR-75-1 and MCF7; and Ewing's sarcomas 6647, RD-ES, and TC-71 were all obtained from the American Type Culture Collection (ATCC, Rockville, MD) and maintained in RPMI + 10% FCS. All cultures tested negative for mycoplasma contamination.

Fresh tumor specimens were prepared from enzymatically dispersed single cell suspensions of solid

- 32 -

- ° tumors, including melanomas, colon carcinomas, sarcomas, and lymphomas. They were cryopreserved in 90% FCS + 10% DMSO, and were rapidly thawed for immediate use on the day of bioassay.

Antigen Presentation to T Cells by EBV-Transformed B Cells

- 5 Optimization of a bioassay for tumor-reactive CD4<sup>+</sup> T cells, using EBV-B cells as antigen presenting cells for lysates of whole tumor cells, has been described (Topalian, S.L., et al. (1994) *Int. J. Cancer*. 58:69-79). Assay medium consisted of RPMI + 10% AB serum with IL-2  
10 120 IU/ml. Briefly, washed EBV-B cells were cultured at 7-10 x 10<sup>5</sup> cells/ml. Tumor antigen was added to B cells in the form of cell lysates: concentrated preparations of washed whole tumor cells were subjected to 3 cycles of rapid freezing and thawing, and cell fragments were added  
15 to B cells at 7-10 x 10<sup>5</sup> cell equivalents/ml. Tumor-pulsed B cell cultures were maintained at 37°C for 20-24 h, then TIL were added at 4-10 x 10<sup>5</sup> cells/ml. Cultures were established in flat-bottom 96-well plates (220 ul/well), 48-well plates (550 ul/well), or 24-well plates  
20 (1100 ul/well) depending on the numbers of cells available. As a positive control, TIL were also cultured in plates coated with anti-CD3 mAb (OKT3; Ortho Pharmaceuticals, Raritan, NJ). Cultures were maintained for an additional 20-24 hours (h), then supernatants were  
25 harvested by centrifugation and stored at -80°C until assayed for the presence of cytokines. Secreted cytokines were measured with ELISA kits purchased from R+D Systems (Minneapolis, MN) for granulocyte-macrophage colony stimulating factor (GM-CSF, detectable concentrations 8-  
30 500 pg/ml), tumor necrosis factor alpha (TNF- $\alpha$ , 15-1000 pg/ml), IL-4 (31-2000 pg/ml) and IL-6 (3-300 pg/ml). GM-CSF assays were calibrated with international reference standard 88/646 (NCI-FCRDC, Frederick, MD). Interferon gamma (IFN- $\gamma$ ) was measured with an ELISA developed with  
35

- 33 -

° reagents from Biosource International (Camarillo, CA) (20-10,000 pg/ml).

The response of TIL to tumor stimulation was considered to be significant when cytokine secretion in response to tumor-pulsed EBV-B cells exceeded the response to EBV-B cells alone by  $\geq 3.0$ -fold.

#### Antibody Blocking Studies.

To inhibit TIL recognition of tumor-pulsed EBV-B cells, stimulator cells were cultured with preservative-free mAb for 30 min at room temperature before adding TIL, and then cultures were maintained for 24 h at 37°C in the continued presence of mAb 23 ug/ml. Antibodies directed against HLA determinants included W6/32 (against HLA-A,B,C; IgG<sub>2a</sub>; Sera-Lab, Sussex, England), IVA12 (HLA-DR, DP, DQ; IgG), L243 (HLA-DR; IgG<sub>2a</sub>), Genox 3.53 and G2b.2 (HLA-DQw1; IgG, and IgG<sub>2a</sub>, respectively), and IVD12 (HLA-DQw3; IgG) (all purified from ATCC hybridoma supernatants).

#### Transfection of COS-7 Cells.

Genes cloned into the expression vectors pCDNA3 (Invitrogen, San Diego, CA) or pCEV27 (Miki, T., et al. (1991) *Proc. Natl. Acad. Sci. USA* **88**, 5167-5171) were transiently transfected into the monkey kidney COS-7 cell line (a gift of Dr. W. Leonard, NIH) using the DEAE dextran method (Seed, B., & Aruffo, A. (1987) *Proc. Natl. Acad. Sci. USA* **84**, 3365-3369). EBV-B cells were pulsed with lysates of transfected COS-7 cells for recognition by T cells. The tyrosinase gene was isolated from a cDNA library from the cultured melanoma line 1290A-mel, and its identity was confirmed by partial DNA sequencing which gave a sequence identical to that published by Bouchard (Bouchard, B., et al. (1989) *J. Exp. Med.* **169**, 2029-2042). The tyrosinase (1-3) (comprising exons 1 through 3) gene, also isolated from the 1290A-mel library, lacks the fourth and fifth exons and encodes a truncated product. The gene encoding tyrosinase related protein (gp75) was isolated by screening a cDNA library from 501-mel with a probe

- 34 -

constructed based on the published gene sequence  
(Vijayasaraadhi, S., et al. (1990) *J. Exp. Med.* 171, 1375-1380).

### RESULTS

CD4<sup>+</sup> T cells were purified by positive selection  
5 from a heterogeneous population of bulk cultured TIL  
derived from a metastatic melanoma lesion from patient  
1088. Selected cultures were >95% CD4<sup>+</sup>. Preliminary  
experiments indicated that these CD4<sup>+</sup> T cells secreted  
cytokines specifically when cocultivated with autologous  
10 EBV-B cells (1088-EBV) which had been pulsed with lysates  
of autologous melanoma cells (1088-mel). TIL secreted  
large quantities of GM-CSF, and much smaller quantities of  
TNF- $\alpha$ , IL-4, and IFN- $\gamma$  in response to autologous tumor;  
specific IL-6 secretion was not observed. Thus, GM-CSF  
15 secretion was monitored as a measure of T cell recognition  
in subsequent assays. In 11 separate experiments, CD4<sup>+</sup>  
TIL stimulated with tumor-pulsed EBV-B cells secreted 8-  
to 138-fold more GM-CSF (median 46-fold) than TIL  
stimulated with EBV-B cells in the absence of tumor. As  
20 shown in Figure 1, GM-CSF secretion could be abrogated by  
blocking with the anti-HLA-DR mAb L243 (96% inhibition),  
suggesting that TIL reactivity was HLA-DR restricted.  
Significant blocking of cytokine secretion was also  
observed by the anti-class II framework mAb IVA12 (71%  
25 inhibition), but not by isotype-matched mAb directed  
against a monomorphic MHC class I determinant or against  
two HLA-DQ determinants (HLA type of patient 1088: HLA-DR  
4, 17; HLA-DQw2, 3; HLA-DRw52, 53). Conversely, in the  
same experiment, cytokine secretion by purified CD8<sup>+</sup> TIL  
30 1088 in response to whole autologous melanoma cells (MHC  
class I<sup>+</sup>, class II<sup>+</sup>) was inhibited by the anti-class I mAb  
W6/32, but not by L243 or IVA12 (data not shown). This  
experiment was repeated once with similar results. CD8<sup>+</sup>  
TIL 1088 capable of recognizing whole 1088-mel cells  
35 failed to react to 1088-EBV cells pulsed with tumor

- 35 -

lylate. These data indicate that CD4<sup>+</sup> TIL 1088 recognize autologous tumor antigen presented by EBV-B cells in a specific, MHC class II-restricted manner.

A variety of allogeneic tumors and normal tissues were screened for the presence of Ag recognized by CD4<sup>+</sup> TIL 1088. A representative experiment is shown in Figure 2, in which lysates of the autologous cultured melanoma line as well as all 7 allogeneic melanoma cultures tested were stimulatory. Stimulation indices [SI = GM-CSF secretion by (TIL + EBV-B + cell lysate)/(TIL + EBV-B)] ranged from 4.2 for 938-mel to 104.8 for 553-mel. TIL alone secreted 22 pg/ml GM-CSF, and when cocultivated with unpulsed 1088-EBV secreted 101 pg/ml, as compared to 10,175 pg/ml secreted in the presence of 1088-EBV pulsed with 1088-mel lysate. In separate experiments, a total of 18 fresh or cultured melanomas were screened for recognition, of which 12 were positive (67%), 4 were negative, and 2 equivocal on repeat experiments. Although tumor 624-mel was recognized by TIL 1088 on 5 separate occasions (SI = 8.0 to 24.8), only 4 of 6 tumor clones from 624-mel were recognized. Taken together, these results suggest that the Ag recognized by CD4<sup>+</sup> TIL 1088 are broadly but not universally expressed in melanoma lesions; alternatively, recognition may reflect relative degrees of antigen expression and the sensitivity of our detection system. Of note, TIL recognized autologous fresh melanoma cells (SI=19.0 and 13.1 in two experiments) as well as tumor cultured from the same lesion (1088-mel), from passages 6 through 48. Thus, the recognized antigen were present in vivo and were not a function of culture artifact; their expression was conserved through almost one year of continuous in vitro culture.

Although TIL recognized a number of allogeneic melanomas on repeat assays, they consistently failed to react with normal cells of nonmelanocyte lineage derived from the same patients. For instance, CD4<sup>+</sup> TIL 1088

- 36 -

° responded to 1088-EBV pulsed with 1088-mel lysate but not to these same EBV-B cells pulsed with 1088-EBV lysate (SI = 35.7 and 1.4, respectively). This experiment was repeated twice with similar results. Also, TIL secreted significant amounts of GM-CSF in response to a lysate of fresh 501B melanoma cells, but not to lysates of EBV-B cells or cultured fibroblasts derived from the same patient (SI = 13.4, 2.2, and 1.5 respectively).

TIL were tested for recognition of a variety of nonmelanoma tumors, both fresh and cultured. As represented in Figure 3, TIL failed to recognize lysates from 14 tumors of various histologic types, including colon carcinomas, breast carcinomas, lymphomas, and sarcomas. TIL also failed to react with two Ewing's sarcomas, which share a neuroectodermal embryonic origin with melanomas and were recognized by melanoma-specific CD8<sup>+</sup> T cells in a previous study (Shamamian, P., et al. (1994) *Cancer Immunol. Immunother.* 39:73-83). However, CD4<sup>+</sup> TIL 1088 did recognize all four normal melanocyte lines assayed, on repeated occasions. Measured levels of GM-CSF secretion approached those observed in response to 1088-mel. These results suggested that the Ag recognized by CD4<sup>+</sup> TIL 1088 might be specific for the melanocytic lineage.

Two other melanoma patients whose CD4<sup>+</sup> TILs recognized lysates of autologous melanoma cells presented by autologous or HLA-matched EBV-B cells had previously been identified (Topalian, S. et al., (1994) *Int. J. Cancer* 58:69-79) by our laboratory. These TILs appeared to be MHC class II-restricted and to recognize antigens unique to autologous melanoma cells, since they failed to react with 15 allogeneic melanomas including 1088-mel or with normal cells including cultured melanocytes. These results suggest the existence of multiple class II-restricted melanoma determinants which are differentially

- 37 -

and specifically recognized by CD4<sup>+</sup> T cells from these three patients.

TIL cultured in bulk under the conditions described have been shown to be oligoclonal, but not monoclonal cell populations (Belldegrun, A., et al. (1989) *J. Immunol.* **142**, 4520-4526; Topalian, S.L., et al. (1990) *J. Immunol.* **144**, 4487-4495; Nishimura, M.I., et al. (1993) *J. Cell. Biochem.* **17D**, 110. (Abstr.)). To determine whether multiple shared antigens were being recognized by CD4<sup>+</sup> TIL 1088, CD4<sup>+</sup> T cell clones were raised from these TIL and assayed for target recognition. As shown in Table 1, 4 clones recognized the autologous melanoma as well as multiple allogeneic melanomas and all three normal melanocyte lines tested (FM 902, 906, 907). The target recognition profiles of all 4 clones were remarkably similar and antibody blocking studies suggested that all were HLA-DR restricted. For 2 clones, HLA-DR restriction was confirmed by using allogeneic EBV-B cell lines or macrophages as APC for tumor antigen; only antigen presenting cells sharing the HLA-DR4 molecule were stimulatory. Thus, a single antigenic protein seemed to be present in all of the melanomas recognized by these CD4<sup>+</sup> T cell clones, and was shared by normal melanocytes. Further experiments with these and 7 additional CD4<sup>+</sup> TIL 1088 clones revealed a homogeneous recognition profile, suggesting the presence of an immunodominant epitope in the 1088 system.

CD4<sup>+</sup> TIL 1088 clones were assessed for recognition of melanoma-associated gene products expressed by 1088-mel on Northern blotting and known to contain commonly expressed CD8 epitopes which can be recognized by CTL derived from melanoma patients. The genes encoding the tyrosinase, MART-1, and gp100 proteins (Brichard, V., et al. (1993) *J. Exp. Med.* **178**, 489-495; Kawakami, Y., et al. (1994) *Proc. Natl. Acad. Sci. USA.* **91**, 3515-3519; Bakker, A.B.H., et al. (1994) *J. Exp. Med.* **179**, 1005-1009;

- 38 -

° Kawakami, Y., et al. (1994) *Proc. Natl. Acad. Sci. USA* 91, 6458-6462) were cloned into plasmid vectors and transiently expressed in monkey kidney COS-7 cells. Lysates of transfected COS-7 cells were pulsed onto 1088-EBV and used to stimulate cytokine secretion from CD4<sup>+</sup> T cell clones. Table 2 shows that all 6 T-cell clones tested secreted significant amounts of GM-CSF in response to lysates of tyrosinase-transfected COS-7 cells as well as to lysates from 1088-mel cells. Transfection of COS-7 cells with a truncated tyrosinase gene, or genes encoding the gp75 tyrosinase-related protein,  $\beta$ -galactosidase, or HLA-A2.1 did not confer recognition. As a control, CD8<sup>+</sup> T cells from patient 1088 failed to react with any of these stimulator cells. Of note, uncloned CD4<sup>+</sup> T cells did not react as strongly as the clones with the tyrosinase gene product. This may reflect the presence of additional epitopes recognized by CD4<sup>+</sup> TIL 1088 subpopulations not represented by these clones, as also suggested by some discrepancies in the recognition of allogeneic melanomas by the bulk-cultured TIL compared to the T cell clones (compare Figure 2 to Table 1). In three additional experiments, CD4<sup>+</sup> T cell clones specifically recognized the products of tyrosinase genes isolated from two different patients' melanomas and expressed in two different plasmids, while failing to react with MART-1 or gp100 (not shown). Taken together with previous demonstrations of CD8<sup>+</sup> T cell reactivity in melanoma patients against HLA-A2 and HLA-A24 restricted epitopes encoded by the tyrosinase gene (Brichard, et al. (1993) *J. Exp. Med.* 178:489-495; Robbins, P.F. et al. (1994) *Cancer Research* 54:3124-3126), these findings show that a single gene product, tyrosinase, contains epitopes recognized by both CD4<sup>+</sup> and CD8<sup>+</sup> T lymphocytes. Furthermore, the fact that the CD4<sup>+</sup> TIL react to lysates of melanomas as well as normal melanocytes suggests that the recognized epitope is nonmutated.



- 39 -

°           The importance of CD4<sup>+</sup> T cells in the priming  
and effector phases of the antitumor immune response has  
been shown in animal models (Greenberg, P.D., et al.  
(1985) *J. Exp. Med.* **161**, 1122-1134; Kern, D.E., et al.  
(1986) *J. Immunol.* **136**, 4303-4310; Ostrand-Rosenberg, S.,  
5 Roby, C.A., & Clements, V.K. (1991) *J. Immunol.* **147**, 2419-  
2422; Dranoff, G., et al. (1993) *Proc. Natl. Acad. Sci.*  
*USA* **90**, 3539-3543). Although experimental immunization  
strategies for patients with melanoma and other cancers  
currently emphasize shared class I restricted tumor  
10 antigens recognized by CD8<sup>+</sup> T cells, immunization against  
both class I and class II restricted epitopes may increase  
the effectiveness of these approaches.

15

20

25

30

35

Table 1. CD4<sup>+</sup> T cell clones recognize a shared antigen expressed on autologous and allogeneic melanomas and normal melanocytes

Stimulator*	GM-CSF secretion, † pg/ml per 24 hr (St†)			
	NT9	ID6	IB7	IE7
1088-mel	14,581 (504)	27,992 (384)	21,497 (169)	13,767 (101)
1011-mel	13,626 (471)	18,937 (260)	15,882 (125)	9,292 (68)
553-mel	2,856 (99)	11,067 (153)	9,357 (74)	4,777 (36)
697-mel	2,451 (86)	8,887 (123)	6,497 (52)	3,382 (26)
526-mel	168 (7)	2,117 (30)	1,400 (12)	196 (2)
624-mel	44 (3)	1,537 (22)	1,187 (10)	98 (2)
1087-mel	37 (2)	806 (12)	675 (6)	53 (1)
1290B-mel	<8 (1)	168 (3)	120 (2)	33 (1)
938-mel	<8 (1)	50 (2)	32 (1)	<8 (1)
1102-mel	<8 (1)	<8 (1)	<8 (1)	<8 (1)
586-mel	<8 (1)	55 (2)	18 (1)	<8 (1)
537-mel	<8 (1)	34 (1)	<8 (1)	<8 (1)
501A-mel	<8 (1)	24 (1)	<8 (1)	<8 (1)
677-mel	<8 (1)	22 (1)	<8 (1)	<8 (1)
CY13	<8 (1)	<8 (1)	<8 (1)	<8 (1)
1088-EBV	<8 (1)	<8 (1)	<8 (1)	<8 (1)
FM 902	5,831 (202)	12,667 (175)	10,517 (83)	6,557 (49)
FM 906	11,376 (393)	17,767 (244)	16,307 (128)	NT
FM 907	9,401 (325)	17,382 (239)	15,202 (120)	NT

NT, not tested.

\* Cell lysates of cultured lines listed were incubated with 1088-EBV cells for 20 hr. In the same experiment, CD8<sup>+</sup> TIL 1088 failed to react with these stimulators (data not shown).† Net secretion = (secretion by TILs with EBV-B plus cell lysate) - (secretion by TILs with EBV-B). All cells were at  $4 \times 10^5$  per ml in microtiter plates.

‡ Stimulation index. GM-CSF secretion by TIL with EBV-B in the absence of cell lysate ranged from 29 to 138 pg/ml.

Table 2. CD4<sup>+</sup> T cell clones recognize a product of the tyrosinase gene

Stimulator*	Transfected gene†	GM-CSF secretion, ‡ pg/ml per 24 hr						
		CD4 <sup>+</sup> bulk TILs	CD8 <sup>+</sup> bulk TILs	NT9	ID6	IB7	IE7	2G2
1088-mel	None	6169	56	32,904	58,515	43,550	17,563	11,702
COS-7	Tyrosinase	123	<8	2,804	7,015	9,150	2,163	12,202
COS-7	Tyrosinase (1-3)	12	<8	<8	14	<8	<8	<8
COS-7	Tyrosinase-related gp75	<8	<8	<8	<8	46	<8	<8
COS-7	β-Galactosidase	8	<8	<8	16	36	16	<8
COS-7	HLA-A2.1	8	<8	<8	22	<8	<8	<8

\* Cell lysates were cocultivated with 1088-EBV for 20 hr.

† Genes were expressed in the plasmid vector pcDNA3 except for the tyrosinase-related gp75 gene, which was in pCEV27.

‡ Net secretion is defined in footnote † to Table 1. Values to TIL with EBV-B ranged from 10 to 237 pg/ml.

- 42 -

## Example II

MHC Class II Restricted Tyrosinase  
Peptides And Modifications ThereofMATERIALS AND METHODS

5        Peptide Synthesis. Peptides were synthesized  
by a solid phase method using a peptide synthesizer, and  
their molecular weights confirmed by mass spectrometry.  
HA 307-319 and MT(65) 3-13, with high affinities for HLA-  
DRB1\*0401 and B1\*0301, respectively, were synthesized for  
10 use as inhibitors of tyrosinase peptide binding (Sette, A.  
et al (1993) *J. Immunol.* 151, 3163-3170; Sidney J. et al  
(1992) *J. Immunol.* 149, 2634-2640, herein incorporated by  
reference).

15        Tumor Cell Lines And Other Cell Lines. Same as  
in Example 1

Isolation of CD4<sup>+</sup> TIL 1088. Same as in Example  
1.

20        Assessment of Antigen Recognition by CD4<sup>+</sup>TIL  
1088. EBV-B cells were incubated overnight in the  
presence of peptides, up to 100 $\mu$ M concentrations. CD4<sup>+</sup> T  
cells were added for an additional 24 hours, and cytokine  
secretion measured to assess T cell stimulation (see  
Example 1).

25        Results. Two tyrosinase peptides recognized by  
bulk CD4<sup>+</sup> TIL1088 have been identified: Ty 56-70 and Ty  
448-462 (Figure 7). This was done by screening  
overlapping 15-mers based on a tyrosinase sequence derived  
from melamona cell line 501-mel, (this sequence is the  
same as GenBank J03581 sequence but with an R instead of Q  
at amino acid position 402). These peptides are  
30 nonmutated, since their amino acid sequences are identical  
to a tyrosinase sequence derived from normal human  
melanocytes (Kwon et. al., *PNAS* (1987) 84:7473-7477;  
GenBank Accession Number J03581).

35        CD4<sup>+</sup> T cells from patient 1088 had been shown to  
recognize lysates of autologous and some allogeneic

- 43 -

° melanoma cell lines expressing tyrosinase, and recognition (measured via cytokine secretion) was HLA-DR-restricted (Example 1; Figures 1 and 2). Patient 1088 was heterozygous for DR, expressing HLA-DRB1\*0301 and B1\*0401. A peptide binding HLA-DRB1\*0401 with high affinity, HA 307-319, effectively blocked binding of both the Ty 56-70 and Ty 448-462 peptides to the presenting EBV-B cells derived from patient 1088, leading to decreased T cell recognition (see Figure 4). A peptide with high affinity for HLA-DRB1\*0301, MT(65)3-13, failed to inhibit T cell recognition. Neither HA 307-319 nor MT(65) 3-13 affected T cell recognition of a melanoma lysate, which is internalized by B cells and processed prior to presentation (Topalian *Int. J. Cancer*, 1994 58:69-79). The HLA-restriction of Ty 56-70 and Ty 448-462 was confirmed as HLA-DRB1\*0401 by using B cell lines of various HLA genotypes as antigen presenting cells (See Figure 5).

In order to make more immunogenic peptides for induction of a CD4<sup>+</sup> T cell response, a variety of peptide epitopes were synthesized in which at least one amino acid position was changed based on the binding motifs of peptides presented by HLA-DRB\*0401 (Sette, A. et al (1993) *J. Immunol.* 151, 3163-3170; Rammensee, H.G. (1995) *Immunogenetics* 41, 178-228, herein incorporated by reference). The P1 anchor position of Ty 56-70 was hypothesized to be I58, L59, or L60 and the P6 anchor to be A63, P64 or L65 (Figure 6). To confirm this, peptides with amino acid substitutions at these positions (Q = unfavorable substitution, F or V = favorable) and serially truncated peptides were tested for T cell recognition. The results shown in Figure 6 suggest that I58 and A63 are the P1 and P6 anchors, respectively, of Ty 56-70 (Figure 7). In addition, two modified peptides (Ty 56-70, A63 → V and Ty 56-70, L65 → V) seemed to evoke an enhanced T cell response compared to the unmodified Ty 56-70. This was

- 44 -

° confirmed by titrating the T cell response to these peptides (Figure 8).

In repeated experiments, Ty 448-462 was specifically recognized by CD4<sup>+</sup> T cells from patient 1088. By testing modified peptides with either unfavorable (Q) or favorable (F) amino acid substitutions at potential P1 anchor positions (Figure 9), and by testing a series of truncated peptides, Y451 was identified as the P1 anchor. At this position, the unfavorable amino acid substitution abolished T cell recognition, while the favorable substitution restored it. In addition, it was found that the truncated Ty 449-462 and 450-462 peptides were more stimulatory for T cells than the parent peptide. This was confirmed in a subsequent titration experiment (see Figure 10).

15 The P6 anchor position for Ty 448-462 was presumed to be D456, based on identifying Y451 as the P1 anchor. D (aspartic acid) is not an optimal residue at this position, and a valine substitution led to markedly enhanced T cell recognition (see Figure 11). When two favorable modifications of Ty 448-462 were combined in a single modified peptide (Ty 450-462, D456→V), recognition was enhanced even more (see Figure 12).

20 In summary, both Ty 56-70 and Ty 448-462 are restricted by HLA-DRB1\*0401. This MHC molecule is expressed by approximately 15% of the North American Caucasian population. The anchor positions in Ty 56-70 and Ty 448-462 have been identified (see Figure 7), and substitutions of amino acids at these positions have created modified peptides with enhanced T cell stimulatory properties.

30 The utility of these peptides in the prophylaxis and/or therapy of melanoma may not be limited to patients expressing the Class II MHC molecule DRB1\*0401, as Class II-restricted peptides are often capable of binding to more than one Class II molecule (Chicz, R.M. et al (1993)

35

- 45 -

- ° J. *Exp. Med.* 178, 27-47; Malcherek, G. et al (1995) *J. Exp. Med.* 181, 527-536).

Although the present invention has been described in some detail by way of illustration, and examples for purposes of clarification and understanding  
5 it will be obvious that certain changes may be made within the scope of the appended claims. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and accompanying  
10 drawings. Such modifications are intended to fall within the scope of the claims.

15

20

25

30

35

- 46 -

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- 5 (i) APPLICANTS: THE GOVERNMENT OF THE UNITED STATES OF AMERICA AS REPRESENTED BY THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES
- (ii) TITLE OF INVENTION: MHC-CLASS II RESTRICTED MELANOMA ANTIGENS AND THEIR USE IN DIAGNOSTIC AND THERAPEUTIC METHODS
- 10 (iii) NUMBER OF SEQUENCES: 38
- (iv) CORRESPONDENCE ADDRESS:  
(A) ADDRESSEE: MORGAN & FINNEGAN, L.L.P.  
(B) STREET: 345 PARK AVENUE  
(C) CITY: NEW YORK  
(D) STATE: NEW YORK  
(E) COUNTRY: USA  
15 (F) ZIP: 10154
- (v) COMPUTER READABLE FORM:  
(A) MEDIUM TYPE: FLOPPY DISK  
(B) COMPUTER: IBM PC COMPATIBLE  
(C) OPERATING SYSTEM: PC-DOS/MS-DOS  
(D) SOFTWARE: ASCII
- 20 (vi) CURRENT APPLICATION DATA:  
(A) APPLICATION NUMBER: TO BE ASSIGNED  
(B) FILING DATE: 25-SEPT-1996  
(C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:  
(A) APPLICATION NUMBER: 08/533,895  
25 (B) FILING DATE: 26 SEPT 1995  
(C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:  
(A) NAME: FEILER, WILLIAM S.  
(B) REGISTRATION NUMBER: 26,728  
(C) REFERENCE/DOCKET NUMBER: 2026-4205PC
- 30 (ix) TELECOMMUNICATION INFORMATION:  
(A) TELEPHONE: (212) 758-4800  
(B) TELEFAX: (212) 751-6849  
(C) TELEX: 421792

## (2) INFORMATION FOR SEQ ID NO:1:

- 35 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 15



- 47 -

(B) TYPE: AMINO ACID  
(C) STRANDEDNESS: UNKNOWN  
(D) TOPOLOGY: UNKNOWN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

5 Gln Asn Ile Leu Leu Ser Asn Ala Pro Leu Gly Pro  
1 5 10

Gln Phe Pro  
15

(2) INFORMATION FOR SEQ ID NO:2:

10

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 14  
(B) TYPE: AMINO ACID  
(C) STRANDEDNESS: UNKNOWN  
(D) TOPOLOGY: UNKNOWN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

15

Asn Ile Leu Leu Ser Asn Ala Pro Leu Gly Pro Gln  
1 5 10

Phe Pro

20 (2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 15  
(B) TYPE: AMINO ACID  
(C) STRANDEDNESS: UNKNOWN  
(D) TOPOLOGY: UNKNOWN

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Gln Asn Ile Leu Leu Ser Asn Ala Pro Val Gly Pro  
1 5 10

Gln Phe Pro  
15

30

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 15  
(B) TYPE: AMINO ACID  
(C) STRANDEDNESS: UNKNOWN  
35 (D) TOPOLOGY: UNKNOWN

- 48 -

o

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Gln Asn Ile Leu Leu Ser Asn Val Pro Val Gly Pro  
 1 5 10

5 Gln Phe Pro  
 15

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

10

(A) LENGTH: 15  
 (B) TYPE: AMINO ACID  
 (C) STRANDEDNESS: UNKNOWN  
 (D) TOPOLOGY: UNKNOWN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

15 Gln Asn Ile Leu Leu Ser Asn Val Pro Leu Gly Pro  
 1 5 10

Gln Phe Pro  
 15

(2) INFORMATION FOR SEQ ID NO:6:

20

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 15  
 (B) TYPE: AMINO ACID  
 (C) STRANDEDNESS: UNKNOWN  
 (D) TOPOLOGY: UNKNOWN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

25

Asp Tyr Ser Tyr Leu Gln Asp Ser Asp Pro Asp Ser  
 1 5 10

Phe Gln Asp  
 15

30

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 15  
 (B) TYPE: AMINO ACID  
 (C) STRANDEDNESS: UNKNOWN  
 (D) TOPOLOGY: UNKNOWN

35

- 49 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Asp Tyr Ser Tyr Leu Gln Asp Ser Asp Pro Asp Ser  
1 5 10

5 Ser Gln Asp  
15

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

10 (A) LENGTH: 15  
(B) TYPE: AMINO ACID  
(C) STRANDEDNESS: UNKNOWN  
(D) TOPOLOGY: UNKNOWN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

15 Asp Gln Ser Tyr Leu Gln Asp Ser Asp Pro Asp Ser  
1 5 10

Phe Gln Asp  
15

(2) INFORMATION FOR SEQ ID NO:9:

20 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 15  
(B) TYPE: AMINO ACID  
(C) STRANDEDNESS: UNKNOWN  
(D) TOPOLOGY: UNKNOWN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

25 Asp Phe Ser Tyr Leu Gln Asp Ser Asp Pro Asp Ser  
1 5 10

Phe Gln Asp  
15

30 (2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 15  
(B) TYPE: AMINO ACID  
(C) STRANDEDNESS: UNKNOWN  
(D) TOPOLOGY: UNKNOWN

35

- 50 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Asp Tyr Ser Phe Leu Gln Asp Ser Asp Pro Asp Ser  
 1 5 10

5 Phe Gln Asp  
 15

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

10 (A) LENGTH: 15  
 (B) TYPE: AMINO ACID  
 (C) STRANDEDNESS: UNKNOWN  
 (D) TOPOLOGY: UNKNOWN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

15 Asp Tyr Ser Tyr Leu Gln Asp Ser Val Pro Asp Ser  
 1 5 10

Phe Gln Asp  
 15

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

20 (A) LENGTH: 13  
 (B) TYPE: AMINO ACID  
 (C) STRANDEDNESS: UNKNOWN  
 (D) TOPOLOGY: UNKNOWN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

25 Ser Tyr Leu Gln Asp Ser Val Pro Asp Ser Phe Gln  
 1 5 10

Asp

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

30 (A) LENGTH: 14  
 (B) TYPE: AMINO ACID  
 (C) STRANDEDNESS: UNKNOWN  
 (D) TOPOLOGY: UNKNOWN

35

- 51 -

° (xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Tyr Ser Tyr Leu Gln Asp Ser Asp Pro Asp Ser Phe  
 1 5 10

Gln Asp

5 (2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 13  
 (B) TYPE: AMINO ACID  
 (C) STRANDEDNESS: UNKNOWN  
 (D) TOPOLOGY: UNKNOWN

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Ser Tyr Leu Gln Asp Ser Asp Pro Asp Ser Phe Gln  
 1 5 10

Asp

15

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 9  
 (B) TYPE: AMINO ACID  
 (C) STRANDEDNESS: UNKNOWN  
 (D) TOPOLOGY: UNKNOWN

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

Xaa Leu Leu Xaa Asn Xaa Xaa Leu Xaa  
 1 5

25 (2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 9  
 (B) TYPE: AMINO ACID  
 (C) STRANDEDNESS: UNKNOWN  
 (D) TOPOLOGY: UNKNOWN

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

Xaa Leu Gln Xaa Ser Xaa Xaa Asp Xaa  
 1 5

35

- 52 -

## ° (2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 15  
(B) TYPE: AMINO ACID  
(C) STRANDEDNESS: UNKNOWN  
(D) TOPOLOGY: UNKNOWN

5

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

Gln Asn Gln Leu Leu Ser Asn Ala Pro Leu Gly Pro  
1 5 10

Gln Phe Pro  
15

10

## (2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 15  
(B) TYPE: AMINO ACID  
(C) STRANDEDNESS: UNKNOWN  
(D) TOPOLOGY: UNKNOWN

15

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

Gln Asn Phe Leu Leu Ser Asn Ala Pro Leu Gly Pro  
1 5 10

20

Gln Phe Pro  
15

## (2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 15  
(B) TYPE: AMINO ACID  
(C) STRANDEDNESS: UNKNOWN  
(D) TOPOLOGY: UNKNOWN

25

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Gln Asn Val Leu Leu Ser Asn Ala Pro Leu Gly Pro  
1 5 10

30

Gln Phe Pro  
15

## (2) INFORMATION FOR SEQ ID NO:20:

35

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 15

- 53 -

(B) TYPE: AMINO ACID  
(C) STRANDEDNESS: UNKNOWN  
(D) TOPOLOGY: UNKNOWN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

5 Gln Asn Ile Gln Leu Ser Asn Ala Pro Leu Gly Pro  
1 5 10  
Gln Phe Pro  
15

(2) INFORMATION FOR SEQ ID NO:21:

10

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 15  
(B) TYPE: AMINO ACID  
(C) STRANDEDNESS: UNKNOWN  
(D) TOPOLOGY: UNKNOWN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

15

Gln Asn Ile Phe Leu Ser Asn Ala Pro Leu Gly Pro  
1 5 10  
Gln Phe Pro  
15

20

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 15  
(B) TYPE: AMINO ACID  
(C) STRANDEDNESS: UNKNOWN  
(D) TOPOLOGY: UNKNOWN

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

Gln Asn Ile Val Leu Ser Asn Ala Pro Leu Gly Pro  
1 5 10  
Gln Phe Pro  
15

30

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 15  
(B) TYPE: AMINO ACID  
(C) STRANDEDNESS: UNKNOWN  
(D) TOPOLOGY: UNKNOWN

35

- 54 -

o

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

Gln Asn Ile Leu Gln Ser Asn Ala Pro Leu Gly Pro  
1 5 10

5 Gln Phe Pro  
15

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

10

(A) LENGTH: 15  
(B) TYPE: AMINO ACID  
(C) STRANDEDNESS: UNKNOWN  
(D) TOPOLOGY: UNKNOWN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

15 Gln Asn Ile Leu Phe Ser Asn Ala Pro Leu Gly Pro  
1 5 10

Gln Phe Pro  
15

(2) INFORMATION FOR SEQ ID NO:25:

20

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 15  
(B) TYPE: AMINO ACID  
(C) STRANDEDNESS: UNKNOWN  
(D) TOPOLOGY: UNKNOWN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

25

Gln Asn Ile Leu Val Ser Asn Ala Pro Leu Gly Pro  
1 5 10

Gln Phe Pro  
15

30 (2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 15  
(B) TYPE: AMINO ACID  
(C) STRANDEDNESS: UNKNOWN  
(D) TOPOLOGY: UNKNOWN

35



- 55 -

° (xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

Gln Asn Ile Leu Leu Ser Asn Gln Pro Leu Gly Pro  
1 5 10  
Gln Phe Pro  
15

5

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 15  
(B) TYPE: AMINO ACID  
(C) STRANDEDNESS: UNKNOWN  
(D) TOPOLOGY: UNKNOWN

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

Gln Asn Ile Leu Leu Ser Asn Ala Gln Leu Gly Pro  
1 5 10  
Gln Phe Pro  
15

15

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 15  
(B) TYPE: AMINO ACID  
(C) STRANDEDNESS: UNKNOWN  
(D) TOPOLOGY: UNKNOWN

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

Gln Asn Ile Leu Leu Ser Asn Ala Val Leu Gly Pro  
1 5 10  
Gln Phe Pro  
15

25

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 15  
(B) TYPE: AMINO ACID  
(C) STRANDEDNESS: UNKNOWN  
(D) TOPOLOGY: UNKNOWN

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

Gln Asn Ile Leu Leu Ser Asn Ala Pro Gln Gly Pro  
1 5 10

35

- 56 -

o

Gln Phe Pro  
15

(2) INFORMATION FOR SEQ ID NO:30:

5

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 13  
(B) TYPE: AMINO ACID  
(C) STRANDEDNESS: UNKNOWN  
(D) TOPOLOGY: UNKNOWN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

10

Ile Leu Leu Ser Asn Ala Pro Leu Gly Pro Gln Phe  
1 5 10

Pro

(2) INFORMATION FOR SEQ ID NO:31:

15

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 13  
(B) TYPE: AMINO ACID  
(C) STRANDEDNESS: UNKNOWN  
(D) TOPOLOGY: UNKNOWN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

20

Leu Leu Ser Asn Ala Pro Leu Gly Pro Gln Phe Pro Phe  
1 5 10

(2) INFORMATION FOR SEQ ID NO:32:

25

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 13  
(B) TYPE: AMINO ACID  
(C) STRANDEDNESS: UNKNOWN  
(D) TOPOLOGY: UNKNOWN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

30

Leu Ser Asn Ala Pro Leu Gly Pro Gln Phe Pro Phe Thr  
1 5 10

(2) INFORMATION FOR SEQ ID NO:33:

35

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 15  
(B) TYPE: AMINO ACID

- 57 -

(C) STRANDEDNESS: UNKNOWN  
(D) TOPOLOGY: UNKNOWN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

5 Ser Asn Ala Pro Leu Gly Pro Gln Phe Pro Phe  
1 5 10  
Thr Gly Val Asp  
15

(2) INFORMATION FOR SEQ ID NO:34:

10 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 15  
(B) TYPE: AMINO ACID  
(C) STRANDEDNESS: UNKNOWN  
(D) TOPOLOGY: UNKNOWN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

15 Asp Tyr Ser Gln Leu Gln Asp Ser Asp Pro Asp Ser  
1 5 10  
Phe Gln Asp  
15

20 (2) INFORMATION FOR SEQ ID NO:35:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 15  
(B) TYPE: AMINO ACID  
(C) STRANDEDNESS: UNKNOWN  
(D) TOPOLOGY: UNKNOWN

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

Asp Tyr Ser Tyr Gln Gln Asp Ser Asp Pro Asp Ser  
1 5 10  
Phe Gln Asp  
15

30

(2) INFORMATION FOR SEQ ID NO:36:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 15  
(B) TYPE: AMINO ACID  
(C) STRANDEDNESS: UNKNOWN  
35 (D) TOPOLOGY: UNKNOWN

- 58 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

Asp Tyr Ser Tyr Phe Gln Asp Ser Asp Pro Asp Ser  
1 5 10

5 Phe Gln Asp  
15

(2) INFORMATION FOR SEQ ID NO:37:

(i) SEQUENCE CHARACTERISTICS:

10

(A) LENGTH: 12  
(B) TYPE: AMINO ACID  
(C) STRANDEDNESS: UNKNOWN  
(D) TOPOLOGY: UNKNOWN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

15 Tyr Leu Gln Asp Ser Asp Pro Asp Ser Phe Gln Asp  
1 5 10

(2) INFORMATION FOR SEQ ID NO:38:

(i) SEQUENCE CHARACTERISTICS:

20

(A) LENGTH: 11  
(B) TYPE: AMINO ACID  
(C) STRANDEDNESS: UNKNOWN  
(D) TOPOLOGY: UNKNOWN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

25 Leu Gln Asp Ser Asp Pro Asp Ser Phe Gln Asp  
1 5 10

30

35

- 59 -

° We Claim:

1. An isolated MHC Class II immunogenic peptide comprising at least 9 contiguous amino acids of a MHC Class II core binding sequence from a tyrosinase sequence.  
5
2. The immunogenic peptide of claim 1 where said peptide contains a sequence selected from the group consisting of QNILLSNAPLGPQFP (Ty 56-70), NILLSNAPLGPQFP (Ty 57-70), DYSYLQSDPDSFQD (Ty 448-462), YSYLQSDPDSFQD  
10 (Ty 449-462), and SYLQSDPDSFQD (Ty 450-462).
3. The immunogenic peptide of claim 2 having at least 9 contiguous residues, wherein said peptide has 1,2, 3, 4 or 5 amino acid deletions at the carboxy or  
15 amino terminus and maintains requisite activity.
4. The immunogenic peptide of claim 1, wherein said peptide sequence contains at least one amino acid modification of said tyrosinase sequence to enhance  
20 binding of peptide to an MHC Class II molecule or to a T cell receptor.
5. The immunogenic peptide of Claim 4 containing a sequence selected from the group consisting  
25 of QNILLSNAPVGPQFP (Ty 56-70, L65→V), QNILLSNVPGPQFP (Ty 56-70, A63→V and L65→V), QNILLSNVPLGPQFP (Ty 56-70, A63→V), DYSYLQSDPDSSQD (Ty 448-462, F460→S), DQSYLQSDPDSFQD (Ty 448-462, Y449→Q), DFSYLQSDPDSFQD (Ty 448-462, Y449→F), DYSFLQSDPDSFQD (Y451→F), and  
30 DYSYLQSDVPDSFQD (Ty 448-462, D456→V).
6. The immunogenic peptide of claim 5 having at least 9 contiguous residues of claim 2, wherein said peptide has 1,2, 3, 4 or 5 amino acid deletions at the  
35 carboxy or amino terminus and maintains requisite activity.

- 60 -

7. The immunogenic peptide of claim 4 wherein an amino acid substitution is within a 9 amino acid Class II binding core sequence in said peptide.

8. The peptide of claim 7, wherein said amino acid substitution is located at a position selected from the group consisting of: (i) the first position, (ii) the fourth position, (iii) the sixth position, (iv) the seventh position, (v) the ninth position and (vi) combinations of at least two of (i) - (v) in the sequence of the core binding region of the peptide.

9. The peptide of claim 6, wherein said amino acid substitution are located at the first and sixth positions.

10. An immunogenic peptide having the core binding sequence formula  $X_1LLX_2NX_3X_4LX_5$  or  $X_1LQX_2SX_3X_4DX_5$ , wherein:

$X_1$  is any hydrophobic amino acid;

$X_2$  is any hydrophobic amino acid; aspartic acid, or glutamic acid;

$X_3$  is any hydrophobic or hydroxyl amino acid;

$X_4$  is any polar, charged or aliphatic amino acid; and

$X_5$  is any polar or aliphatic amino acid.

11. The peptide of claim 10 wherein the peptide is lengthened by flanking regions at the amino and/or carboxy termini to a total of 34 amino acids.

12. The peptide of claim 10, wherein  $X_1$  is selected from the group consisting of methionine, leucine, isoleucine, tyrosine, valine, tryptophan and phenylalanine.

- 61 -

°           13. The peptide of claim 10, wherein  $X_2$  is selected from the group consisting of phenylalanine, tryptophan, leucine, isoleucine, alanine, valine, aspartic acid and glutamic acid.

5           14. The peptide of claim 10, wherein  $X_3$  is selected from the group consisting of methionine, leucine, threonine, isoleucine, serine and valine.

10           15. The peptide of claim 10, wherein  $X_4$  is selected from the group consisting of aspartic acid, alanine, serine, valine, histidine, proline, asparagine, methionine, threonine, leucine and isoleucine.

15           16. The peptide of claim 10, wherein  $X_5$  is selected from the group consisting of alanine, serine, glutamine, glycine, leucine, valine and threonine.

20           17. The immunogenic peptide of claim 1 wherein said peptide is recognized by HLA-DR CD4<sup>+</sup> T lymphocytes.

          18. The immunogenic peptide of claim 10 wherein said peptide is recognized by HLA-DR CD4<sup>+</sup> T lymphocytes.

25           19. The immunogenic peptide of claim 1 linked to a MHC Class II molecule.

          20. The immunogenic peptide of claim 10 linked to a MHC Class II molecule.

30           21. The immunogenic peptide-MHC complex of claim 19 wherein said MHC Class II molecule is the  $\beta$  chain of the MHC Class II molecule.

35           22. The immunogenic peptide-MHC complex of claim 20 wherein said MHC Class II molecule is the  $\beta$  chain of the MHC Class II molecule.

- 62 -

23. A pharmaceutical composition comprising at least one peptide of claim 1, and an acceptable excipient, diluent or carrier.

24. A pharmaceutical composition comprising at least one peptide of claim 10, and an acceptable excipient, diluent or carrier.

25. A pharmaceutical composition comprising at least one peptide of claim 19, and an acceptable excipient, diluent or carrier.

26. A pharmaceutical composition comprising at least one peptide of claim 20, and an acceptable excipient, diluent or carrier.

27. A method of preventing or treating melanoma comprising administering a therapeutically effective amount of the the pharmaceutical composition of claim 23 to a mammal in an effective amount to stimulate the production of protective antibodies or immune cells.

28. A method of preventing or treating melanoma comprising administering a therapeutically effective amount of the the pharmaceutical composition of claim 24 to a mammal in an effective amount to stimulate the production of protective antibodies or immune cells.

29. A method of preventing or treating melanoma comprising administering a therapeutically effective amount of the the pharmaceutical composition of claim 25 to a mammal in an effective amount to stimulate the production of protective antibodies or immune cells.

30. A method of preventing or treating melanoma comprising administering a therapeutically effective amount of the the pharmaceutical composition of claim 26



- 63 -

° to a mammal in an effective amount to stimulate the production of protective antibodies or immune cells.

31. A purified and isolated nucleic acid sequence encoding a peptide according to claim 1.

5

32. A purified and isolated nucleic acid sequence encoding a peptide according to claim 10.

33. A purified and isolated nucleic acid sequence encoding a peptide according to claim 19.

10

34. A purified and isolated nucleic acid sequence encoding a peptide or polypeptide according to claim 20.

15

35. A recombinant expression vector comprising at least one nucleic acid sequence of claim 1.

36. A recombinant expression vector comprising at least one nucleic acid sequence claim 10.

20

37. A recombinant expression vector comprising at least one nucleic acid sequence of claim 19.

38. A recombinant expression vector comprising at least one nucleic acid sequence of claim 20.

25

39. The vector of claim 35, wherein the expression vector is a eukaryotic expression vector or prokaryotic expression vector.

30

40. The vector of claim 36, wherein the expression vector is a eukaryotic expression vector or prokaryotic expression vector.

35

- 64 -

°           41. The vector of claim 37, wherein the  
expression vector is a eukaryotic expression vector or  
prokaryotic expression vector.

5           42. The vector of claim 38, wherein the  
expression vector is a eukaryotic expression vector or  
prokaryotic expression vector.

10           43. A host cell containing with the recombinant  
expression vector according to claim 39.

          44. A host cell containing with the recombinant  
expression vector according to claim 40.

15           45. A host cell containing with the recombinant  
expression vector according to claim 41.

          46. A host cell containing with the recombinant  
expression vector according to claim 42.

20           47. Antibodies reactive with an immunogenic  
peptide of claim 1.

          48. Antibodies reactive with an immunogenic  
peptide of claim 10.

25           49. Antibodies reactive with an immunogenic  
peptide of claim 19.

30           50. Antibodies reactive with an immunogenic  
peptide of claim 20.

          51. The antibodies of claim 48 wherein said  
antibodies are monoclonal.

35           52. A therapeutic method of preventing or  
treating melanoma comprising administering the

- 65 -

- ° pharmaceutical composition of claim 23 to a mammal in an effective amount to stimulate the production of protective antibodies or immune positive CD4<sup>+</sup> T-cells.

5 53. A therapeutic method of preventing or treating melanoma comprising administering the pharmaceutical composition of claim 24 to a mammal in an effective amount to stimulate the production of protective antibodies or immune positive CD4<sup>+</sup> T-cells.

10 54. A therapeutic method of preventing or treating melanoma comprising administering the pharmaceutical composition of claim 25 to a mammal in an effective amount to stimulate the production of protective antibodies or immune positive CD4<sup>+</sup> T-cells.

15 55. A therapeutic method of preventing or treating melanoma comprising administering the pharmaceutical composition of claim 26 to a mammal in an effective amount to stimulate the production of protective  
20 antibodies or immune positive CD4<sup>+</sup> T-cells.

56. The method of claim 52, wherein said composition includes isolated tyrosine protein.

25 57. A method for determining or isolating Class II tumor associated antigens, said method comprising the steps of:

(a) contacting a candidate antigen with antigen presenting cells for a time sufficient to allow  
30 the antigen to be processed by said antigen presenting cells;

(b) contacting said antigen presenting cells step (a) with CD4<sup>+</sup> T lymphocytes; and

(c) screening for recognition of said  
35 antigen presenting cells by said CD4<sup>+</sup> T lymphocytes.

- 66 -

58. The method of claim 57 wherein said candidate antigen is selected from the group consisting of a crude cellular lysate, a purified protein, a peptide, or proteins encoding by a DNA library.

59. The method of claim 57 wherein said antigen presenting cells are EBV transformed B cells, monocytes or dendritic cells.

60. The method of claim 57 wherein said candidate antigen is selected from the group consisting of MART-1, gp100, gp-75, MAGE-1, MAGE-3, or p15 or any other tumor associated protein known to be recognized by CD8<sup>+</sup>T cells via MHC Class I restriction.

61. The melanoma antigens or tumor associated antigens isolated by screening candidate antigens according to claim 57.

62. A purified and isolated nucleic acid sequence encoding a peptide comprising at least about 9 contiguous amino acids, said peptide being derived from a tyrosinase sequence, said peptide being reactive to CD4<sup>+</sup> T-lymphocytes.

63. A recombinant expression vector comprising at least one nucleic acid sequence of claim 62.

64. The use of the peptides of claims 1-22 in the manufacture of a medicament for stimulating the production of protective antibodies or immune cells in a mammal.

65. The use of the peptides of claims 1-22 in the manufacture of a medicament for preventing or treating melanoma by stimulating the production of protective antibodies or immune positive CD4<sup>+</sup> T-cells.




FIGURE 1




FIGURE 2




FIGURE 3

CD4+ TIL 1088 RECOGNIZE TY 56-70 AND  
TY 448-462 PRESENTED BY HLA-DR4




FIGURE 4



## HLA Restriction of Tyrosinase Recognition

by CD4<sup>+</sup> TIL 1088

(GM-CSF secretion, pg/ml/24 hr)

Antigen Presenting Cells	Antigen Presented				HLA-DR81*	
	None	1088-mal lysate	Ty 56-70 (A63-V)	Ty 448-462 (D456-V)		
<u>Experiment 1</u>						
1088-EBV	145	1179	2445	2969	0301	0401
697-EBV	282	1014	>500	3815	0401	1501
EM14	751	3282	4793	5121	0401	-
860-EBV	376	870	383	345	0301	1301
583-EBV	484	550	574	477	0404	1501
1087-EBV	507	557	550	566	0701	1201
<u>Experiment 2</u>						
1359-EBV	183	2238	1179	1714	0301	0401
EM14	372	2822	1892	2805	0401	-
1485-EBV	171	160	154	222	0407	1501
COX	544	482	735	703	0301	-
PGF	54	150	76	46	1501	-

Antigen presenting cells pulsed with tumor lysate or 100  $\mu$ M peptide overnight,  
then washed prior to assay, all cells at  $1 \times 10^6$ /ml

FIGURE 5

# Determining the P1 and P6 Anchor Positions for Ty 56-70

		GM-CSF (pg/ml/24h.)		
		Exp. 1	Exp. 2	Exp. 3
Ty 56-70	Q N I L L S N A P L G P Q F P	295	524	903
I58	- - Q - - - - - - - - -	<8	<8	<8
	- - F - - - - - - - - -	39	48	ND
	- - V - - - - - - - - -	ND	ND	154
L59	- - - Q - - - - - - - - -	<8	<8	<8
	- - - F - - - - - - - - -	<8	<8	ND
	- - - V - - - - - - - - -	ND	ND	<8
L60	- - - - Q - - - - - - - - -	<8	62	<8
	- - - - F - - - - - - - - -	88	<8	ND
	- - - - V - - - - - - - - -	ND	ND	10
A63	- - - - - - - Q - - - - - -	<8	<8	ND
	- - - - - - - V - - - - - -	889	2080	ND
P64	- - - - - - - Q - - - - - -	<8	<8	ND
	- - - - - - - V - - - - - -	<8	48	ND
L65	- - - - - - - Q - - - - - -	24	50	ND
	- - - - - - - V - - - - - -	2050	3816	ND
Ty 57-70	- - - - - - - - - - - - - -	ND	ND	615
Ty 58-70	- - - - - - - - - - - - - -	ND	<8	<8
Ty 59-71	- - - - - - - - - - - - - -	ND	<8	<8
Ty 60-72	- - - - - - - - - - - - - -	ND	<8	<8
Ty 61-75	- - - - - - - - - - - - - -	ND	<8	ND
1088mel Lysate		6762	>5500	ND

TIL & EBV backgrounds for Exp.1=53, Exp.2=296, Exp.3=99 pg/ml.  
 These have been subtracted from the numbers shown above.

FIGURE 6

Ty 56-70

Q N I L L S N A P L G P Q F P

Ty 448-462

D Y S Y L Q D S D P D S F Q D

FIGURE 7

# CD4+ 1088 TIL RECOGNIZE MODIFIED TYROSINASE 56-70 PEPTIDES




FIGURE 8

# Determining the P1 Anchor for Ty 448-462

		GM-CSF (pg/ml/24h.)	
		<u>Exp. 1</u>	<u>Exp. 2</u>
Ty 448-462	D Y S Y L Q D S D P D S F Q D	108	802
Y449	- Q - - - - - - - - - -	74	520
	- F - - - - - - - - - -	103	781
Y451	- - - Q - - - - - - - - -	28	<8
	- - - F - - - - - - - - -	324	1065
L452	- - - - Q - - - - - - - - -	14	<8
	- - - - F - - - - - - - - -	14	<8
Ty 449-462	- - - - - - - - - - - - -	485	1960
Ty 450-462	- - - - - - - - - - - - -	556	2749
Ty 451-462	- - - - - - - - - - - - -	21	<8
Ty 452-462	- - - - - - - - - - - - -	18	<8
1088-mel Lysate		>5500	5822

TIL & EBV backgrounds for Exp.1=52 and Exp.2=196 pg/ml.  
 These have been subtracted from the numbers shown above.

FIGURE 9

### CD4+ TIL 1088 RECOGNIZE MODIFIED TYROSINASE 448-462 PEPTIDES






FIGURE 10

**CD4+ TIL 1088 Recognize Modified Ty 448-462****FIGURE 11**

# CD4+ TIL 1088 RECOGNIZE MODIFIED TYROSINASE 448 - 462 PEPTIDES




FIGURE 12